

Arithmetic on Elliptic Threefolds

Rania Wazir

February 1, 2008

1 Introduction

Consider an elliptic curve $E/\mathbb{Q}$ given by the Weierstrass equation

$$E : y^2 = x^3 + Ax + B; \quad \text{with } A, B \in \mathbb{Z}, \quad (1)$$

and with discriminant locus

$$\Delta := 4A^3 + 27B^2 \neq 0.$$

The Mordell-Weil Theorem shows that $E(\mathbb{Q})$, the set of rational points on E , is a finitely-generated Abelian group.

If in equation (1) we instead let the coefficients A, B lie in a polynomial ring over $\mathbb{Z}$, then E is no longer defined over $\mathbb{Q}$, but over some function field K of $\mathbb{Q}$, and we obtain an **elliptic fibration** or **elliptic n-fold**. The function-field analogue of the Mordell-Weil Theorem shows that, in this case also, the rational points on E are a finitely-generated Abelian group. The rank of $E(K)$ has been an object of intense study and speculation, yet many of its properties, and the relation to the underlying geometry of E , remain elusive. The aim of this paper is to prove a relation between the rank of an elliptic threefold, and an average of its fibral Frobenius trace values.

Theorem 1.1 *Let k be a number field, $\mathfrak{p}$ a prime in k , and $q_{\mathfrak{p}}$ its norm.*

Let $\mathcal{E} \rightarrow \mathcal{S}$ be a non-split elliptic threefold defined over k . Then Tate's Conjecture for $\mathcal{E}/k$ and $\mathcal{S}/k$ implies

$$\operatorname{res}_{s=1} \sum_{\mathfrak{p}} -A_{\mathfrak{p}}(\mathcal{E}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} = \operatorname{rank} \mathcal{E}(\mathcal{S}/k).$$

1.1 History of the problem

Consider an elliptic curve E over $\mathbb{Q}(T)$. This is known as an **elliptic surface**, and has Weierstrass equation of the form:

$$E : y^2 = x^3 + A(T)x + B(T); \quad \text{with } A(T), B(T) \in \mathbb{Z}[T],$$

with discriminant locus

$$\Delta(T) := 4A(T)^3 + 27B(T)^2 \neq 0.$$

For each $t \in \mathbb{Z}$ and each prime p , let

$$a_p(E_t) := 1 + p - \#\{\text{rational points in the reduction of } E_t \bmod p\}$$

and let

$$A_p(E) := \frac{1}{p} \sum_{t=1}^p a_p(E_t)$$

be the average of the a_p 's over all fibers.

Then, based on calculations of some non-trivial examples, Nagao [22] conjectured that

$$\lim_{X \rightarrow \infty} \frac{1}{X} \sum_{p \leq X} -A_p(E) \log p = \text{rank } E(\mathbb{Q}(T)).$$

Recently, Rosen and Silverman [25] have derived an analytic version of Nagao's formula:

$$\text{res}_{s=1} \sum_p -A_p(E) \frac{\log p}{p^s} = \text{rank } E(\mathbb{Q}(T)).$$

Assuming Tate's Conjecture, they prove that the analytic version of Nagao's formula holds for elliptic surfaces defined over any number field k , and, with a mild non-vanishing assumption, that in fact the original Nagao formula holds.

Theorem 1.1 gives an analytic Nagao formula for elliptic threefolds.

1.2 Outline of the Proof

The proof proceeds along the following lines:

1. We first prove an isomorphism in the cohomology of $\mathcal{E}$ and $\mathcal{S}$:

$$H_{\text{ét}}^1(\mathcal{S}/\bar{k}, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\mathcal{E}/\bar{k}, \mathbb{Q}_l)$$

as $\text{Gal}(\bar{k}/k)$ -modules.

2. Next, we need a Shioda-Tate formula for elliptic threefolds.
3. Find a geometric interpretation of the Frobenius action on the singular fibers of $\mathcal{E} \rightarrow \mathcal{S}$.

4. The number of rational points on a fibered variety over $\mathbb{F}_{\mathfrak{p}}$ can be counted in two ways: we can count the number of rational points on the threefold (using the Lefschetz Fixed Point Formula); or we can count the number of rational points fiber by fiber, and take the sum over all fibers. Thus, taking the reduction $\tilde{\mathcal{E}}$ of $\mathcal{E} \bmod (\mathfrak{p})$, and equating the two expressions for the number of rational points on $\tilde{\mathcal{E}}$ will give an equality involving $A_{\mathfrak{p}}(\mathcal{E})$.
5. Re-interpret the equation for $A_{\mathfrak{p}}(\mathcal{E})$ in terms of L -series, and apply Tate's Conjecture.
6. Plug in the results from steps 1, 2, and 3.

Remark 1.1 *Steps 1 and 2 are actually proven in the more general case of an elliptic n -fold. The need for an effective geometric Tchebotarev theorem requires restriction to the elliptic threefolds case for the remaining Steps.*

Remark 1.2 *The results in this article formed the bulk of the author's Ph.D. thesis [32].*

2 Basic Definitions and Notation

Most of the results given in this paper, except for the section on singular fibers, and the concluding section on L -series, apply in higher dimensions, and not just in the case of elliptic threefolds. Therefore we give here the general definition of an elliptic n -fold, and where necessary, restrict to the case of an elliptic threefold. This section also contains other basic definitions and auxiliary results that will be needed in the rest of the article. For convenience, a list of notation used is included at the end of the paper.

Notation 2.1 *Let k be a number field, with ring of integers O_k , and algebraic closure $\bar{k}$.*

2.1 Elliptic N-Folds

Definition 2.1 *An elliptic n -fold defined over k is a smooth, projective variety $\mathcal{E}/k$ of dimension n , together with a proper, flat k -morphism $\pi : \mathcal{E} \rightarrow \mathcal{S}$ to a smooth projective $(n-1)$ -dimensional variety $\mathcal{S}/k$, such that the generic fiber is a smooth elliptic curve E defined over $k(\mathcal{S})$, the function field of $\mathcal{S}/k$.*

In general, *morphism* means a morphism defined over $\bar{k}$, and *k -morphism* means a morphism defined over k .

Definition 2.2 *A section $\sigma : \mathcal{S} \rightarrow \mathcal{E}$ is a morphism such that the composition $\pi \circ \sigma = \text{id}_{\mathcal{S}/\bar{k}}$. A **k -section** $\sigma : \mathcal{S} \rightarrow \mathcal{E}$ is a k -morphism such that the composition $\pi \circ \sigma = \text{id}_{\mathcal{S}/k}$.*

Definition 2.3 A rational section $\sigma : \mathcal{S} \rightarrow \mathcal{E}$ is a rational map (defined over $\bar{k}$) such that the composition $\pi \circ \sigma = \text{id}_{\mathcal{U}/\bar{k}}$, for some affine open subset $\mathcal{U} \subset \mathcal{S}$.

A **k-rational section** $\sigma : \mathcal{S} \rightarrow \mathcal{E}$ is a rational map (defined over k) such that the composition $\pi \circ \sigma = \text{id}_{\mathcal{U}/k}$, for some affine open subset $\mathcal{U} \subset \mathcal{S}$.

Definition 2.4 An elliptic n -fold $\mathcal{E} \rightarrow \mathcal{S}$ defined over k is **split** (over $\bar{k}$) if there is an elliptic curve $\mathfrak{E}$ defined over $\bar{k}$, and a birational isomorphism (over $\bar{k}$) $\mu : \mathcal{E} \xrightarrow{\sim} \mathfrak{E} \times_{\bar{k}} \mathcal{S}$, such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{\mu} & \mathfrak{E} \times_{\bar{k}} \mathcal{S} \\ \pi \searrow & & \swarrow \text{proj}_2 \\ & \mathcal{S} & \end{array}$$

Notation 2.2 Define $\mathcal{E}/k$ to be a non-split elliptic n -fold $\pi : \mathcal{E} \rightarrow \mathcal{S}$ defined over k , with k -section $\sigma_0 : \mathcal{S} \rightarrow \mathcal{E}$. Assume also that $\mathcal{E}(k) \neq \emptyset$.

Remark 2.1 Note that, because of the section σ_0 , $\mathcal{E}(k) \neq \emptyset$ is equivalent to $\mathcal{S}(k) \neq \emptyset$. This assumption is made in order to ensure that $\text{Pic}_{\mathcal{E}}^0$ and $\text{Pic}_{\mathcal{S}}^0$ are defined over k . (See the discussion in [12], pp. 31-33).

Notation 2.3 Denote by (O) the image of the section σ_0 in $\mathcal{E}$, and by O the corresponding rational point on $\mathcal{S}$.

Definition 2.5 The closed subset

$$\Delta := \{s \in \mathcal{S} \mid \mathcal{E}_s \text{ is not regular}\}$$

is called the **discriminant locus** of $\mathcal{E}$. Δ is a divisor on $\mathcal{S}$. Let $\Delta = \Delta_1 + \cdots + \Delta_r$ be the irreducible decomposition of Δ .

Notation 2.4

$$\begin{aligned} K &= k(\mathcal{S}), \text{ the function field of } \mathcal{S}/k. \\ \bar{K} &\text{ the algebraic closure of } K. \\ \widehat{K} &= \bar{k}(\mathcal{S}), \text{ the function field of } \mathcal{S}/\bar{k}. \\ \mathcal{O}_{\mathcal{S}} &\text{ the structure sheaf of } \mathcal{S}/\bar{k}. \\ \mathcal{O}_{\mathcal{E}} &\text{ the structure sheaf of } \mathcal{E}/\bar{k}. \end{aligned}$$

We now make a few observations regarding the elliptic n -fold $\mathcal{E}$.

Proposition 2.1

- (a) $\pi_*(\mathcal{O}_{\mathcal{E}}) \cong \mathcal{O}_{\mathcal{S}}$.
- (b) The fibers $\mathcal{E}_s$ are connected.
- (c) The requirement that the morphism π be flat is equivalent to requiring that $\dim(\mathcal{E}_s)$ be constant for all $s \in \mathcal{S}$.

Proof.

- (a) This follows essentially from the properness of the map π , and the existence of a global section σ .
 1. The morphism $\pi : \mathcal{E} \rightarrow \mathcal{S}$ can be factored into $\pi = q \circ \pi'$, where $\mathcal{S}' := \mathbf{Spec} \pi_* \mathcal{O}_{\mathcal{E}}$, $\pi' : \mathcal{E} \rightarrow \mathcal{S}'$ is a proper morphism with $\pi'_* \mathcal{O}_{\mathcal{E}} = \mathcal{O}_{\mathcal{S}'}$, and $q : \mathcal{S}' \rightarrow \mathcal{S}$ is a finite morphism. ([11], proof of Corollary III.11.5).
 2. By construction, $\mathcal{S}'$ is connected because $\mathcal{E}$ and $\mathcal{S}$ are. Furthermore, the section $\sigma : \mathcal{S} \rightarrow \mathcal{E}$ induces a section $\tilde{\sigma} : \mathcal{S} \rightarrow \mathcal{S}'$, given by $\tilde{\sigma} := \pi' \circ \sigma$. Thus, since $\tilde{\sigma}$ maps $\mathcal{S}$ isomorphically onto its image $\tilde{\sigma}(\mathcal{S}) \subset \mathcal{S}'$, $\dim(\mathcal{S}) = \dim(\tilde{\sigma}(\mathcal{S}))$. But since the morphism q is finite, $\dim(\mathcal{S}) = \dim(\mathcal{S}')$ also. This implies that $\mathcal{S}$ is isomorphic to a connected component of $\mathcal{S}'$, and, since $\mathcal{S}'$ is connected, the result follows:

$$\mathcal{S} \cong \tilde{\sigma}(\mathcal{S}) \cong \mathcal{S}'.$$

- (b) This follows from part (a), and ([11], III.11.3).
- (c) EGA ([10], IV.15.4.2).

♠

2.2 The $\widehat{K}/\bar{k}$ -trace of E

In order to state the Mordell-Weil Theorem for function fields, it is necessary to recall first the concept of the $\widehat{K}/\bar{k}$ -trace of E .

Let $\mathfrak{k}$ be a field, and let $\mathfrak{F}$ be a finitely generated extension of $\mathfrak{k}$, such that $\mathfrak{F}$ is the function field of a variety defined over $\mathfrak{k}$. Let $\mathfrak{A}$ be an Abelian variety defined over $\mathfrak{k}$. An $\mathfrak{F}/\mathfrak{k}$ -trace of $\mathfrak{A}$ is a pair $(\mathfrak{t}, \mathfrak{B})$, consisting of an Abelian variety $\mathfrak{B}$, and a homomorphism $\mathfrak{t} : \mathfrak{B} \rightarrow \mathfrak{A}$ defined over $\mathfrak{F}$, such that $(\mathfrak{t}, \mathfrak{B})$ satisfies the following universal mapping property:

If $(\mathfrak{r}, \mathfrak{C})$ is another pair with $\mathfrak{C}$ defined over $\mathfrak{k}$, and $\mathfrak{r}$ a homomorphism $\mathfrak{r} : \mathfrak{C} \rightarrow \mathfrak{A}$ defined over $\mathfrak{F}$, then there exists a unique homomorphism $\mathfrak{i} : \mathfrak{C} \rightarrow \mathfrak{B}$ defined over $\mathfrak{F}$ making the following diagram commute:

$$\begin{array}{ccc}
& & \mathfrak{C} \\
& \swarrow i & \downarrow \tau \\
\mathfrak{B} & \xrightarrow{\mathfrak{t}} & \mathfrak{A}
\end{array}$$

Chow defined and proved the existence of the $\mathfrak{F}/\mathfrak{k}$ -trace, and also that the homomorphism $\mathfrak{t}$ is injective. See ([12], p.138) for details.

Let (τ, B) be the $\widehat{K}/\bar{k}$ -trace of E . Next, we show that the assumption that the elliptic n -fold $\mathcal{E}$ is non-split implies that $B = 0$.

Lemma 2.1 *Let $\mathcal{E}$, E be as previously defined, and assume that the $\widehat{K}/\bar{k}$ -trace (τ, B) of E is such that $B \neq 0$. Then $\mathcal{E}$ is split over $\bar{k}$.*

Proof. Since $B \neq 0$, B is an Abelian variety of dimension at least one. Since also the morphism τ is injective, we get

$$1 \leq \dim B \leq \dim E = 1;$$

hence $\dim B = \dim E = 1$. Therefore the morphism $\tau : B \rightarrow E$ is an injective isogeny, and so

$$\tau : B \xrightarrow{\sim} E \quad \text{is an isomorphism over } \widehat{K}.$$

This in turn implies $\widehat{K}(B) \cong \widehat{K}(E)$ as $\widehat{K}$ -algebras. Since

$$\bar{k}(B \times_{\bar{k}} \mathcal{S}) = \widehat{K}(B) \cong \widehat{K}(E) = \bar{k}(\mathcal{E}),$$

this induces a birational isomorphism of varieties

$$\mathcal{E} \rightarrow B \times_{\bar{k}} \mathcal{S}$$

commuting with maps to $\mathcal{S}$; by definition, this means $\mathcal{E}$ is split. ♠

2.3 k -Rational Sections

Let $\mathcal{E}(\mathcal{S}/\bar{k})$ be the set of rational sections $\rho : \mathcal{S} \rightarrow \mathcal{E}$, and $\mathcal{E}(\mathcal{S}/k)$ be the set of k -rational sections of $\mathcal{E}$. $\mathcal{E}(\mathcal{S}/\bar{k})$ and $\mathcal{E}(\mathcal{S}/k)$ are groups, and are isomorphic to $E(\widehat{K})$ and $E(K)$ respectively. Proofs and further details are exactly as in the elliptic surfaces case, see ([29], III.3.10).

Definition 2.6 $\mathcal{E}(\mathcal{S}/k)$, the group of $\mathbf{k}$ -rational sections on $\mathcal{E}$, is often called the **Mordell-Weil group** of $\mathcal{E}$.

The next theorem, known as the **Mordell-Weil Theorem for function fields**, will show that both $\mathcal{E}(\mathcal{S}/k)$ and $\mathcal{E}(\mathcal{S}/\bar{k})$ have finite rank.

Theorem 2.1 (Néron - Lang) [14] *Let $\mathfrak{k}$ be a field, and $\mathfrak{K}$ a function field over $\mathfrak{k}$. Let A be an Abelian variety defined over $\mathfrak{K}$, with $\mathfrak{K}/\mathfrak{k}$ -trace $(\mathfrak{t}, \mathfrak{B})$. Then the group of rational points $A(\mathfrak{K})$ modulo the subgroup $\mathfrak{t}\mathfrak{B}(\mathfrak{k})$ is a finitely generated Abelian group.*

Since $B = 0$ by Lemma 2.1, this theorem shows that

$$E(\widehat{K})/\tau B(\bar{k}) = E(\widehat{K}) \cong \mathcal{E}(\mathcal{S}/\bar{k})$$

is finitely generated, and therefore $\text{rank } E(\widehat{K}) = \text{rank } \mathcal{E}(\mathcal{S}/\bar{k})$ is finite. Furthermore, $\mathcal{E}(\mathcal{S}/k)$ is a subgroup of $\mathcal{E}(\mathcal{S}/\bar{k})$, so $\mathcal{E}(\mathcal{S}/k)$ is also finitely generated; $\text{rank } \mathcal{E}(\mathcal{S}/k)$ is often called the **Mordell-Weil rank** of $\mathcal{E}$.

Remark 2.2 *Actually, $\mathcal{E}(\mathcal{S}/k)$ is finitely generated even when $\mathcal{E}(\mathcal{S}/\bar{k})$ is not; this follows from Theorem 2.1, because the Mordell-Weil Theorem for number fields shows that $B(k)$ is finitely-generated.*

2.4 Fibral Frobenius Trace Values

Having defined one side of the equation in Theorem 1 (the Mordell-Weil rank of $\mathcal{E}$), we now address the terms on the other side of the equation, and make more precise our notion of “average” of fibral Frobenius trace values:

Notation 2.5

$\mathbb{F}_{\mathfrak{p}}$	the residue field of a prime $\mathfrak{p}$ of O_k .
$\bar{\mathbb{F}}_{\mathfrak{p}}$	the algebraic closure of $\mathbb{F}_{\mathfrak{p}}$.
$q_{\mathfrak{p}}$	the norm of $\mathfrak{p}$, i.e., $q_{\mathfrak{p}} = \#\mathbb{F}_{\mathfrak{p}}$.
$\text{Frob}_{\mathfrak{p}}$	the Frobenius morphism over $\mathbb{F}_{\mathfrak{p}}$.

For a given smooth, projective variety $\mathcal{V}/k$, denote by $\tilde{\mathcal{V}}$ its reduction mod($\mathfrak{p}$). $\tilde{\mathcal{V}}/\bar{\mathbb{F}}_{\mathfrak{p}}$ will be used to denote the variety $\tilde{\mathcal{V}} \times_{\bar{\mathbb{F}}_{\mathfrak{p}}} \bar{\mathbb{F}}_{\mathfrak{p}}$.

Some of the traces in the Lefschetz Fixed-Point Theorem occur frequently in this paper, so we make the following definitions:

Definition 2.7 *For any smooth, projective variety $\mathcal{V}/k$ and any $\mathfrak{p}$ such that $\tilde{\mathcal{V}}/\bar{\mathbb{F}}_{\mathfrak{p}}$ is smooth, let:*

$$\begin{aligned} a_{\mathfrak{p}}(\mathcal{V}) &:= \text{Trace}(\text{Frob}_{\mathfrak{p}}|H_{\text{ét}}^1(\tilde{\mathcal{V}}/\bar{\mathbb{F}}_{\mathfrak{p}}, \mathbb{Q}_l)), \\ b_{\mathfrak{p}}(\mathcal{V}) &:= \text{Trace}(\text{Frob}_{\mathfrak{p}}|H_{\text{ét}}^2(\tilde{\mathcal{V}}/\bar{\mathbb{F}}_{\mathfrak{p}}, \mathbb{Q}_l)), \\ c_{\mathfrak{p}}(\mathcal{V}) &:= \text{Trace}(\text{Frob}_{\mathfrak{p}}|H_{\text{ét}}^3(\tilde{\mathcal{V}}/\bar{\mathbb{F}}_{\mathfrak{p}}, \mathbb{Q}_l)). \end{aligned}$$

Returning to the case of our elliptic n -fold $\mathcal{E}$, we make the following definitions:

Definition 2.8 For a given point $x \in \tilde{S}(\mathbb{F}_{\mathfrak{p}})$, let

$$a_{\mathfrak{p}}(\mathcal{E}_x) := 1 - \#\tilde{\mathcal{E}}_x(\mathbb{F}_{\mathfrak{p}}) + q_{\mathfrak{p}}m_x,$$

where m_x is the number of $\mathbb{F}_{\mathfrak{p}}$ -rational components of the fiber $\tilde{\mathcal{E}}_x$.

These $a_{\mathfrak{p}}(\mathcal{E}_x)$ will be called the **fibral Frobenius trace values** of $\tilde{\mathcal{E}}$.

Remark 2.3 Notice that by the Lefschetz Fixed-Point Theorem, when the fiber $\tilde{\mathcal{E}}_x$ is smooth, definition 2.11 agrees with definition 2.10 above.

And finally, the “average” of fibral Frobenius trace values:

Definition 2.9

$$A_{\mathfrak{p}}(\mathcal{E}) := \frac{1}{q_{\mathfrak{p}}^{(n-1)}} \sum_{x \in \tilde{S}(\mathbb{F}_{\mathfrak{p}})} a_{\mathfrak{p}}(\tilde{\mathcal{E}}_x).$$

2.5 Integral Models

Consider an embedding of $\mathcal{E}$ into projective space $\mathbb{P}^N$. Since $\mathcal{E}$ is defined over k , we can find a finite collection $\{f_1, \dots, f_t\}$ of homogeneous polynomials, with coefficients $\{A_I\}$ in k , that define $\mathcal{E}$. Singularities on $\mathcal{E}$ are determined by the simultaneous vanishing of these polynomials, and some combination of products of their partial derivatives (again finitely many, again with coefficients $\{B_J\}$ in k). Elimination Theory then states that there is a set of polynomials $\{g_h\}_1^u$ in the coefficients $\{A_I, B_J\}$, such that this is equivalent to the simultaneous vanishing of the g_h . Since $\mathcal{E}$ is nonsingular, these g_h do not simultaneously vanish, and hence there are only finitely many primes $\mathfrak{p}$ such that reduction of the $\{A_I\}$ and $\{B_J\} \bmod(\mathfrak{p})$ would give vanishing of the g_h . Call these the “bad primes”, and collect them in a set B . The complement of B is an open set $U \subset O_k$; $\tilde{\mathcal{E}}$, the reduction of $\mathcal{E} \bmod(\mathfrak{p})$, is still a smooth, elliptic n -fold (defined over $\mathbb{F}_{\mathfrak{p}}$) for all $\mathfrak{p} \in U$, and so the model

$$\begin{array}{c} \mathcal{E} \\ \downarrow \\ \text{Spec } k \end{array}$$

can be extended to an integral model

$$\begin{array}{c} \mathcal{E}_U \\ \downarrow \\ U. \end{array}$$

$\mathcal{E}/k$ can be thought of as the fiber over the generic point of this model, while $\tilde{\mathcal{E}}$ is the fiber over the closed point $\mathfrak{p} \in U$.

Similar Elimination Theory arguments can be used to show that the following hold for all but finitely many primes $\mathfrak{p}$:

1. $\tilde{\mathcal{S}}$ is a smooth $(n - 1)$ -fold (defined over $\mathbb{F}_{\mathfrak{p}}$).
2. $\tilde{\pi} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{S}}$ is a proper, flat morphism.
3. If Δ' is the discriminant locus of $\tilde{\pi} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{S}}$, then for almost all $\mathfrak{p}$, $\Delta' = \tilde{\Delta}$, the reduction mod($\mathfrak{p}$) of Δ , and the irreducible decomposition of Δ' is given by:

$$\Delta' = \tilde{\Delta}_1 + \cdots + \tilde{\Delta}_r.$$

4. Let $\Theta_{i,j}$ be the irreducible components of $\pi^{-1}(\Delta_j)$, and $\Theta'_{i,j}$ be the irreducible components of $\tilde{\pi}^{-1}(\tilde{\Delta}_j)$. Then $\tilde{\Theta}_{i,j} = \Theta'_{i,j}$.
5. For a fixed $x \in \mathcal{S}$, the number of irreducible components in $\tilde{\mathcal{E}}_x$ is the same as the number of irreducible components in $\mathcal{E}_x$.

By enlarging B if necessary, we will assume that these statements hold for all $\mathfrak{p} \in U$.

2.6 Intersection Theory

One of the main results of the next section will be an isomorphism of the Picard Varieties of $\mathcal{E}$ and $\mathcal{S}$. In the case of elliptic surfaces, the proof relies on a non-degenerate bilinear pairing

$$\mathrm{Div}(\mathcal{E}) \times \mathrm{Div}(\mathcal{E}) \rightarrow \mathbb{Z}$$

given by Intersection Theory on surfaces. (Details can be found in [29] Proposition III.8.2). In the case of higher-dimensional varieties, it is no longer possible to get a pairing into $\mathbb{Z}$. However, we will show that it suffices to have a pairing with a notion of “positivity.” This pairing will be the subject of this section. For a review of the necessary results on Intersection Theory of higher-dimensional varieties, see [11], Appendix A or [7]. For ease of reference, we list here only the most necessary results.

For the rest of this section, assume all varieties are non-singular, projective over an algebraically closed field κ .

2.6.1 Summary of Basic Results

Definition 2.10 A cycle $\Gamma = \sum a_i [V_i]$ is **non-negative** if $a_i \geq 0$ for all i ; in that case, we write $\Gamma \geq 0$. Γ is **positive**, written $\Gamma > 0$, if in addition, $a_i > 0$ for some i .

Definition 2.11 Γ is **non-positive**, written $\Gamma \leq 0$, if $a_i \leq 0$ for all i . Γ is **negative**, written $\Gamma < 0$, if in addition, $a_i < 0$ for some i .

Definition 2.12 A cycle class $\alpha \in A_r(X)$ is **non-negative**, written $\alpha \geq 0$, if there is a non-negative r -cycle Γ such that $\Gamma \geq 0$ and $cl(\Gamma) = \alpha$.

Similarly for α **positive** ($\alpha > 0$), α **non-positive** ($\alpha \leq 0$), and α **negative** ($\alpha < 0$).

Theorem 2.2 If $f : X \rightarrow W$ is a proper morphism, and Γ is an r -cycle on W which is algebraically equivalent to zero, then $f_*\Gamma$ is algebraically equivalent to zero on X .

Proof. See Fulton([7], Proposition 10.3). ♠

Theorem 2.3 If $f : X \rightarrow W$ is a morphism, and Γ is an r -cycle on W which is algebraically equivalent to zero, then $f^*\Gamma$ is algebraically equivalent to zero on X .

Proof. See Fulton([7], Example 19.3.9, p.390). ♠

2.6.2 A Nondegenerate Pairing

Definition 2.13 Define a symmetric, bilinear pairing

$$\langle \cdot, \cdot \rangle : \text{Pic}(\mathcal{E}) \times \text{Pic}(\mathcal{E}) \rightarrow \text{Pic}(\mathcal{S})$$

via

$$\langle \Lambda, \Upsilon \rangle := \pi_*(\Lambda \cdot \Upsilon)$$

for any $\Lambda, \Upsilon \in \text{Pic}(\mathcal{E})$.

If C, D are divisors in $\text{Div}(\mathcal{E})$, let $\langle C, D \rangle := \langle cl(C), cl(D) \rangle$.

The following Proposition is very similar to the Elliptic Surfaces case. However, there are enough differences due to the new pairing, that we give a proof below.

Proposition 2.2 Let $D \in \text{Div}(\mathcal{E})$ be a fibral divisor, and $G \in \text{Div}(\mathcal{S})$. Then

- (a) $\langle D, \pi^*(G) \rangle = 0$
- (b) $\langle D, D \rangle \leq 0$
- (c) If $\langle D, D \rangle = 0$ then $D \in \pi^*(\text{Div}(\mathcal{S}))$.

Proof.

- (a) This follows from the Projection Formula, once we note that for any fibral divisor D , $\pi_*(D) = 0$.

$$\begin{aligned}
\langle D, \pi^*(G) \rangle &= \pi_*(cl(D).cl(\pi^*G)) \\
&= \pi_*(D.\pi^*(G)) \\
&= \pi_*(D).G \\
&= 0.G \\
&= 0.
\end{aligned}$$

- (b) Write

$$D = D_1 + D_2 + \dots + D_n,$$

where each D_i is contained in a different fiber. Then $\langle D_i, D_j \rangle = 0$ for $i \neq j$, which implies that

$$\langle D, D \rangle = \langle D_1, D_1 \rangle + \langle D_2, D_2 \rangle + \dots + \langle D_n, D_n \rangle.$$

Thus, it suffices to prove the theorem for each D_i , and we can assume that $D \subset \pi^*G$, for some irreducible $G \in \text{Div}(S)$.

Let

$$F := \pi^*G = \sum_{i=0}^t n_i \Gamma_i$$

be the irreducible decomposition of F . Notice that $n_i \geq 0$ for all i . Furthermore, since $D \subset F$, D can be written as

$$D = \sum_{i=0}^t a_i \Gamma_i.$$

Rewrite D as

$$D = \sum_{i=0}^t \left(\frac{a_i}{n_i}\right) n_i \Gamma_i,$$

and define another fibral divisor D' by

$$D' = \sum_{i=0}^t \left(\frac{a_i}{n_i}\right)^2 n_i \Gamma_i.$$

By part (a), we know that $\langle D', F \rangle = 0$ and thus:

$$\begin{aligned}
-2\langle D, D \rangle &= \langle D', F \rangle - 2\langle D, D \rangle + \langle F, D' \rangle \\
&= \sum_{i,j=0}^t \frac{a_i^2}{n_i^2} \langle n_i \Gamma_i, n_j \Gamma_j \rangle - 2 \sum_{i,j=0}^t \frac{a_i a_j}{n_i n_j} \langle n_i \Gamma_i, n_j \Gamma_j \rangle \\
&\quad + \sum_{i,j=0}^t \frac{a_j^2}{n_j^2} \langle n_i \Gamma_i, n_j \Gamma_j \rangle \\
&= \sum_{i,j=0}^t \left(\frac{a_i}{n_i} - \frac{a_j}{n_j}\right)^2 \langle n_i \Gamma_i, n_j \Gamma_j \rangle
\end{aligned}$$

The terms in the above equation are zero when $i = j$; it therefore reduces to:

$$\langle D, D \rangle = -\frac{1}{2} \sum_{\substack{i,j=0 \\ i \neq j}}^t \left(\frac{a_i}{n_i} - \frac{a_j}{n_j} \right)^2 \langle n_i \Gamma_i, n_j \Gamma_j \rangle \quad (2)$$

Now for $i \neq j$, Γ_i and Γ_j are distinct irreducible divisors, and so

$$\langle \Gamma_i, \Gamma_j \rangle \geq 0 \quad \text{for all } i \neq j.$$

Since, as previously mentioned, $n_i, n_j \geq 0$, this implies that

$$\langle n_i \Gamma_i, n_j \Gamma_j \rangle \geq 0 \quad \text{for all } i \neq j,$$

and therefore $\langle D, D \rangle \leq 0$, proving part (b).

(c) To prove part (c), now assume that $\langle D, D \rangle = 0$. Plugging into equation (2), this gives

$$\frac{a_i}{n_i} = \frac{a_j}{n_j} \quad \text{for all } i, j \text{ such that } \langle \Gamma_i, \Gamma_j \rangle > 0.$$

However, the fibers of π are connected by Proposition 2.1.b. Thus, given any two irreducible components Γ_i and Γ_j , there is a sequence of components

$$\Gamma_i = \Gamma_{i_0}, \Gamma_{i_1}, \dots, \Gamma_{i_r} = \Gamma_j$$

such that Γ_{i_k} has non-empty intersection with $\Gamma_{i_{k+1}}$. Hence

$$\frac{a_i}{n_i} = \frac{a_j}{n_j} \quad \text{for all } i, j.$$

But the section $\sigma : S \rightarrow \mathcal{E}$ intersects with exactly one irreducible component of π^*G , say Γ_0 , which therefore has multiplicity one: i.e., $n_0 = 1$. This implies

$$\frac{a_i}{n_i} = a_0 \in \mathbb{Z} \quad \text{for all } i.$$

Plugging this into the irreducible decomposition of D gives:

$$\begin{aligned} D &= \sum_{i=0}^t a_i \Gamma_i \\ &= \sum_{i=0}^t \frac{a_i}{n_i} n_i \Gamma_i \\ &= \sum_{i=0}^t a_0 n_i \Gamma_i \\ &= a_0 \sum_{i=0}^t n_i \Gamma_i \\ &= a_0 F \in \pi^*(\text{Div}(\mathcal{S})), \end{aligned}$$

This completes the proof of part (c).



2.7 Picard Varieties

We will adopt the following notation for varieties X/κ , κ an algebraically closed field:

Notation 2.6

Pic_X	<i>the Picard Scheme of X/κ.</i>
$\text{Pic}(X)$	$= \text{Pic}_X(\bar{\kappa})$, <i>the Picard Group of $X/\bar{\kappa}$.</i>
$\text{Pic}(X/\kappa)$	$= \text{Pic}_X(\kappa)$, <i>the Picard Group of X/k.</i>
Pic_X^0	<i>the Picard Variety of X/κ.</i>
$\text{Pic}^0(X)$	$= \text{Pic}_X^0(\bar{\kappa})$.
$\text{Pic}^0(X/\kappa)$	$= \text{Pic}_X^0(\kappa)$.
$\text{NS}(X)$	$= \text{Pic}(X)/\text{Pic}^0(X)$, <i>the Neron-Severi Group of $X/\bar{\kappa}$.</i>
$\text{NS}(X/\kappa)$	$= \text{Pic}(X/\kappa)/\text{Pic}^0(X/\kappa)$, <i>the Neron-Severi Group of X/κ.</i>

Definition 2.14 *The trivial part of $\text{NS}(\mathcal{E}) \otimes \mathbb{Q}$, denoted $\mathcal{T}$, is the subspace generated by the image of the zero section (O) , and by all geometrically irreducible components of the fibral divisors.*

Denote by $\mathcal{F}$ the subspace of $\mathcal{T}$ generated by the non-identity components of the fibral divisors, where the identity component of a fibral divisor is the component intersecting (O) .

Note that $\mathcal{T}$ is generated by (O) , $\pi^*\text{NS}(\mathcal{S})$, and $\mathcal{F}$.

Remark 2.4 *For all but finitely many primes $\mathfrak{p}$, the trivial part of $\text{NS}(\tilde{\mathcal{E}}/\bar{\mathbb{F}}_{\mathfrak{p}}) \otimes \mathbb{Q}$ is isomorphic to $\tilde{\mathcal{T}}$, the reduction of $\mathcal{T} \bmod (\mathfrak{p})$. This follows from Section 2.5, statement 3, and hence holds for all primes $\mathfrak{p} \in U$.*

3 An Isomorphism in Cohomology

Our main goal in this section will be to prove

Theorem 3.1

$$H_{\text{ét}}^1(\mathcal{S}/\bar{k}, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\mathcal{E}/\bar{k}, \mathbb{Q}_l).$$

as $\text{Gal}(\bar{k}/k)$ -modules.

This can be proven by showing first that $\text{Pic}_{\mathcal{E}}^0 \cong \text{Pic}_{\mathcal{S}}^0$, and combining this with the fact that, for any variety $\mathcal{V}/k$, $H_{\text{ét}}^1(\mathcal{V}/k, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\text{Pic}_{\mathcal{V}}^0/\bar{k}, \mathbb{Q}_l)$.

The isomorphism of Picard Varieties will follow quite easily from the following theorem:

Theorem 3.2 *There is an exact sequence of Abelian Groups:*

$$0 \longrightarrow \mathrm{Pic}^0(\mathcal{S}) \longrightarrow \mathrm{Pic}^0(\mathcal{E}) \longrightarrow B(\bar{k}). \quad (3)$$

Proof. This proof is based on Raynaud's proof of the elliptic surface case (see [28], Theorem 2).

1. The morphism $\pi : \mathcal{E} \rightarrow \mathcal{S}$ induces a map

$$\pi^* : \mathrm{Pic}_{\mathcal{S}} \rightarrow \mathrm{Pic}_{\mathcal{E}}.$$

Since π^* preserves algebraic equivalence (Theorem 2.3), we get a morphism

$$\alpha := \pi^*|_{\mathrm{Pic}_{\mathcal{S}}^0} : \mathrm{Pic}_{\mathcal{S}}^0 \rightarrow \mathrm{Pic}_{\mathcal{E}}^0.$$

Furthermore, restriction to the generic fiber induces a morphism

$$\psi : \mathrm{Pic}_{\mathcal{E}}^0 \times_{\mathcal{S}} \mathcal{S} \rightarrow \mathrm{Pic}_{\mathcal{E}}^0 \cong E. \quad (4)$$

But by the universal mapping property of the $\widehat{K}/\bar{k}$ -trace (τ, B) of E , the map ψ factors through B ; i.e., there is a unique homomorphism $\beta : \mathrm{Pic}_{\mathcal{E}}^0 \rightarrow B$ such that $\psi = \tau \circ \beta$. Thus, we have a sequence of morphisms

$$\mathrm{Pic}_{\mathcal{S}}^0 \xrightarrow{\alpha} \mathrm{Pic}_{\mathcal{E}}^0 \xrightarrow{\beta} B \quad (5)$$

and it remains to show that, as maps on the $\bar{k}$ -points, this is a short exact sequence of Abelian groups:

$$0 \longrightarrow \mathrm{Pic}^0(\mathcal{S}) \xrightarrow{\alpha} \mathrm{Pic}^0(\mathcal{E}) \xrightarrow{\beta} B(\bar{k}). \quad (6)$$

2. The injectivity of α follows from the existence of the global section $\sigma : \mathcal{S} \rightarrow \mathcal{E}$;

Since by definition $\pi \circ \sigma = id_{\mathcal{S}}$, this gives

$$id_{\mathrm{Pic}(\mathcal{S})} = (\pi \circ \sigma)^* = \sigma^* \circ \pi^*;$$

therefore π^* , and hence also α , is injective.

3. To show exactness at the middle, note first that $\psi \circ \alpha = 0$; this is true because restriction to the generic fiber sends fibral divisors in $\mathrm{Pic}^0(\mathcal{E})$ to zero, and α sends $\mathrm{Pic}^0(\mathcal{S})$ to fibral divisors in $\mathrm{Pic}^0(\mathcal{E})$.

But if $\psi \circ \alpha = 0$, then also $\beta \circ \alpha = 0$ because τ is injective.

This shows $\mathrm{Im}(\alpha) \subset \mathrm{Ker}(\beta)$.

4. Finally, to show $\text{Ker}(\beta) = \text{Im}(\alpha)$, take any $0 \neq \Gamma \in \text{Ker}(\beta)$. Then $\Gamma = cl(D)$ for some divisor D with $D|_E = 0$, and thus D must be a fibral divisor. Since $cl(D) \in \text{Pic}^0(\mathcal{E})$, it follows that $\langle D, F \rangle = 0$ for every divisor F on $\mathcal{E}$, and in particular, $\langle D, D \rangle = 0$. By Theorem (2.2.c), this implies $D \in \pi^*(\text{Div}(\mathcal{S}))$.

Thus, we have $D \sim_{alg} 0$, and $D = \pi^*C$ for some $C \in \text{Div}(\mathcal{S})$. It remains to show that $C \sim_{alg} 0$. But by Theorem 2.3, σ^* preserves algebraic equivalence. Therefore, $\sigma^*D \sim_{alg} 0$. Since

$$\begin{aligned} \sigma^*D &= \sigma^*\pi^*C \\ &= (\pi \circ \sigma)^*C \\ &= (id_{\text{Pic}(\mathcal{S})})C \\ &= C, \end{aligned}$$

this shows that $C \sim_{alg} 0$, and therefore $\Gamma = cl(D) \in \pi^*(\text{Pic}^0(\mathcal{S}))$, proving that $\text{Ker}(\beta) = \text{Im}(\alpha)$.

♠

Now we are ready to prove the isomorphism of Picard Varieties:

Theorem 3.3 $\text{Pic}_{\mathcal{S}}^0$ and $\text{Pic}_{\mathcal{E}}^0$ are isomorphic as Abelian Varieties over k .

Proof.

1. Since the elliptic threefold $\mathcal{E}$ is non-split, the $\widehat{K}/\bar{k}$ -trace B must be trivial (Lemma 2.1). From the exact sequence (3), it then follows that $\text{Pic}^0(\mathcal{S})$ and $\text{Pic}^0(\mathcal{E})$ are isomorphic as groups.
2. If $\text{Pic}^0(\mathcal{S}) \cong \text{Pic}^0(\mathcal{E})$, then $\text{Pic}_{\mathcal{S}}^0 \cong \text{Pic}_{\mathcal{E}}^0$.

For any abelian variety A/k , and any integer l , let

$$A_l := \{a \in A \mid la = 0\}.$$

Then by [19], Proposition II.6, p.64,

$$A_l \cong \mathbb{Z}/l\mathbb{Z}^{2d_A},$$

where $d_A := \dim(A)$.

Therefore, $\text{Pic}^0(\mathcal{S}) \cong \text{Pic}^0(\mathcal{E})$ implies $\text{Pic}^0(\mathcal{S})_l \cong \text{Pic}^0(\mathcal{E})_l$, and so

$$\dim(\text{Pic}^0(\mathcal{S})) = \dim(\text{Pic}^0(\mathcal{E})).$$

Since α is a map between two Abelian varieties of the same dimension, and with finite (in fact, trivial) kernel, this implies that α is an isogeny. ([16], Proposition 8.1.c)

But then

$$\text{Pic}_{\mathcal{E}}^0 \cong \text{Pic}_{\mathcal{S}}^0 / \text{Ker}(\alpha), \quad \text{by [19], Corollary III.10.1, p.118,}$$

and since $\ker(\alpha) = 0$, this gives $\text{Pic}_{\mathcal{E}}^0 \cong \text{Pic}_{\mathcal{S}}^0$.

3. Since Pic_S^0 , Pic_E^0 , and the map α are all defined over k , the isomorphism is in fact an isomorphism over k .

♠

Proposition 3.1 *Let $\mathcal{V}$ be a smooth, projective variety defined over k . Then*

$$H_{\text{ét}}^1(\mathcal{V}/\bar{k}, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\text{Pic}_{\mathcal{V}}^0/\bar{k}, \mathbb{Q}_l).$$

Proof. This seems to be a well-known result, but since we could not find a reference, we give a proof below.

Denote by $\text{Alb}_{\mathcal{V}}$ the Albanese variety of $\mathcal{V}$, and let

$$\mu : \mathcal{V} \longrightarrow \text{Alb}_{\mathcal{V}}$$

be the Albanese map of $\mathcal{V}$. This map is defined over k if $\mathcal{V}(k)$ is not empty - ([13], pp.31-32). Then μ induces an isomorphism in cohomology ([9], p.331)

$$H_1(\mathcal{V}, \mathbb{Z})/\mathfrak{T} \xrightarrow{\sim} H_1(\text{Alb}_{\mathcal{V}}, \mathbb{Z}),$$

where $\mathfrak{T}$ is the torsion subgroup of $H_1(\mathcal{V}, \mathbb{Z})$. Rewrite this as a short exact sequence

$$0 \longrightarrow \mathfrak{T} \longrightarrow H_1(\mathcal{V}, \mathbb{Z}) \longrightarrow H_1(\text{Alb}_{\mathcal{V}}, \mathbb{Z}) \longrightarrow 0.$$

Taking $\text{Hom}(\cdot, \mathbb{Z}/l^r\mathbb{Z})$ of the sequence gives:

$$0 \longrightarrow \text{Hom}(H_1(\text{Alb}_{\mathcal{V}}, \mathbb{Z}), \mathbb{Z}/l^r\mathbb{Z}) \longrightarrow \text{Hom}(H_1(\mathcal{V}, \mathbb{Z}), \mathbb{Z}/l^r\mathbb{Z}) \longrightarrow \text{Hom}(\mathfrak{T}, \mathbb{Z}/l^r\mathbb{Z}).$$

Since $\text{Hom}(H_1(X, \mathbb{Z}), G) \cong H^1(X, G)$ for any finite Abelian group G and any smooth variety X , we get

$$0 \longrightarrow H^1(\text{Alb}(\mathcal{V}), \mathbb{Z}/l^r\mathbb{Z}) \longrightarrow H^1(\mathcal{V}, \mathbb{Z}/l^r\mathbb{Z}) \longrightarrow \text{Hom}(\mathfrak{T}, \mathbb{Z}/l^r\mathbb{Z}).$$

Furthermore, by the étale cohomology compatibility theorems ([15], Theorem III.3.12, p.117), we have $H^1(X, G) \cong H_{\text{ét}}^1(X, G)$ for any finite Abelian group G and any smooth variety X/k :

$$0 \longrightarrow H_{\text{ét}}^1(\text{Alb}_{\mathcal{V}}, \mathbb{Z}/l^r\mathbb{Z}) \longrightarrow H_{\text{ét}}^1(\mathcal{V}, \mathbb{Z}/l^r\mathbb{Z}) \longrightarrow \text{Hom}(\mathfrak{T}, \mathbb{Z}/l^r\mathbb{Z}).$$

Taking inverse limits as r goes to infinity, and noting that inverse limits of exact sequences of finite groups are exact ([15], p.165):

$$0 \longrightarrow H_{\text{ét}}^1(\text{Alb}_{\mathcal{V}}, \mathbb{Z}_l) \longrightarrow H_{\text{ét}}^1(\mathcal{V}, \mathbb{Z}_l) \longrightarrow \varprojlim \text{Hom}(\mathfrak{T}, \mathbb{Z}/l^r\mathbb{Z}),$$

where by definition,

$$H_{\text{ét}}^1(\mathcal{V}, \mathbb{Z}_l) := \varprojlim H_{\text{ét}}^1(\mathcal{V}, \mathbb{Z}/l^r\mathbb{Z}).$$

Furthermore, since $\mathcal{V}$ is a smooth variety, $H_1(\mathcal{V}, \mathbb{Z})$, and hence also $\mathfrak{T}$, is finitely-generated. Therefore, there is some integer t such that $\text{Hom}(\mathfrak{T}, \mathbb{Z}/l^r\mathbb{Z}) \cong \mathbb{Z}/l^t\mathbb{Z}$ for all $r \geq t$, and therefore

$$\varprojlim \text{Hom}(\mathfrak{T}, \mathbb{Z}/l^r\mathbb{Z}) \cong \mathbb{Z}/l^t\mathbb{Z}.$$

But $\mathbb{Z}/l^t\mathbb{Z} \otimes \mathbb{Q} = 0$, hence tensoring with $\mathbb{Q}$ gives

$$0 \longrightarrow H_{\text{ét}}^1(\text{Alb}_{\mathcal{V}}, \mathbb{Z}_l) \otimes \mathbb{Q} \longrightarrow H_{\text{ét}}^1(\mathcal{V}, \mathbb{Z}_l) \otimes \mathbb{Q} \longrightarrow 0.$$

Since $H_{\text{ét}}^1(\mathcal{V}, \mathbb{Q}_l) := H_{\text{ét}}^1(\mathcal{V}, \mathbb{Z}_l) \otimes \mathbb{Q}$, this shows that

$$H_{\text{ét}}^1(\mathcal{V}, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\text{Alb}_{\mathcal{V}}, \mathbb{Q}_l).$$

Noting now that there is an isomorphism defined over k

$$\text{Alb}(\text{Pic}_{\mathcal{V}}^0) \cong \text{Alb}_{\mathcal{V}}, \quad [12], \text{ Thm VI.1.1, p. 148}$$

we see that

$$\begin{aligned} H_{\text{ét}}^1(\text{Pic}_{\mathcal{V}}^0, \mathbb{Q}_l) &\cong H_{\text{ét}}^1(\text{Alb}(\text{Pic}_{\mathcal{V}}^0), \mathbb{Q}_l) \\ &\cong H_{\text{ét}}^1(\text{Alb}_{\mathcal{V}}, \mathbb{Q}_l) \\ &\cong H_{\text{ét}}^1(\mathcal{V}, \mathbb{Q}_l). \end{aligned}$$

as desired. ♠

This leads us to the main theorem of this section:

Theorem 3.1

$$H_{\text{ét}}^1(\mathcal{S}/\bar{k}, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\mathcal{E}/\bar{k}, \mathbb{Q}_l).$$

as $\text{Gal}(\bar{k}/k)$ -modules.

Proof. This follows from Theorem 3.3, together with Proposition 3.1. ♠

The following Corollary will be especially useful in later simplifying the equation for $A_{\mathfrak{p}}(\mathcal{E})$.

Corollary 3.1 $a_{\mathfrak{p}}(\mathcal{E}) = a_{\mathfrak{p}}(\mathcal{S})$.

Proof. . Since Theorem 3.1 gives

$$H_{\text{ét}}^1(\mathcal{E}, \mathbb{Q}_l) \cong H_{\text{ét}}^1(\mathcal{S}, \mathbb{Q}_l),$$

it is clear that

$$\text{Trace}(\text{Frob}_{\mathfrak{p}} | H_{\text{ét}}^1(\mathcal{E}/\bar{k}, \mathbb{Q}_l)) = \text{Trace}(\text{Frob}_{\mathfrak{p}} | H_{\text{ét}}^1(\mathcal{S}/\bar{k}, \mathbb{Q}_l)).$$

Thus, what remains is to show, for any variety $\mathcal{V}/k$, that

$$\text{Trace}(\text{Frob}_{\mathfrak{p}} | H_{\text{ét}}^1(\mathcal{V}/\bar{k}, \mathbb{Q}_l)) = \text{Trace}(\text{Frob}_{\mathfrak{p}} | H_{\text{ét}}^1(\tilde{\mathcal{V}}/\bar{\mathbb{F}}_{\mathfrak{p}}, \mathbb{Q}_l)),$$

and this is a consequence of the proper and smooth base change theorems (see [1]). ♠

4 A Shioda-Tate Formula

In this section, we prove that $\text{NS}(\mathcal{E}) \otimes \mathbb{Q}$ is generated by $\mathcal{T}$ and by the k -rational sections. In the case of elliptic surfaces, this is the main result of the **Shioda-Tate Formula**, which we recall here:

Theorem 4.1 (Shioda-Tate Formula) ([27], Theorem 1.1)

Let $\mathfrak{E} \rightarrow \mathfrak{C}$ be an elliptic surface defined over k , with k -rational section σ_0 .

Embed $\mathfrak{E}(\mathfrak{C}/\bar{k})$ into $\text{NS}(\mathfrak{E})$ by sending a section σ to the divisor $\sigma(\mathfrak{C}) - \sigma_0(\mathfrak{C})$. Then there is a decomposition of $\text{Gal}(\bar{k}/k)$ -modules,

$$\text{NS}(\mathfrak{E}) \otimes \mathbb{Q} \cong (\mathfrak{E}(\mathfrak{C}/\bar{k}) \otimes \mathbb{Q}) \oplus \mathcal{J},$$

where $\mathcal{J}$ is the subspace of $\text{NS}(\mathfrak{E}) \otimes \mathbb{Q}$ generated by the image of the zero section, and by all components of all fibers.

In order to prove a similar formula for elliptic n -folds, we follow an argument similar to Shioda's proof for Elliptic Surfaces [28], and take a closer look at the map ψ (equation 4, restriction to the generic fiber) at the level of geometric points.

Restricting a line bundle on $\mathcal{E}$ to the generic fiber E defines a homomorphism

$$\text{Pic}(\mathcal{E}) \rightarrow \text{Pic}(E/\widehat{K}) \quad (7)$$

which associates with every divisor class $\text{cl}(D)$ on $\mathcal{E}$ the divisor $D|_E = D.E$ on the generic fiber E . Then, using the given rational point $O \in E(K)$, adjust the image by sending $\text{cl}(D)$ to $\text{cl}(D')$, where $D' := D.E - (D.E)O$; the divisor D' is thus a degree zero divisor on E , and the homomorphism becomes

$$\phi : \text{Pic}(\mathcal{E}) \rightarrow \text{Pic}^0(E/\widehat{K}) \cong E(\widehat{K}). \quad (8)$$

Remark 4.1 *By construction, the kernel of ϕ contains the zero-section and any fibral divisor in $\text{Pic}(\mathcal{E})$.*

Lemma 4.1 *Let $\check{\mathcal{T}}$ be the subgroup of $\text{Pic}(\mathcal{E})$ generated by the irreducible components of the fibral divisors, and by the zero-section (O) . Then*

$$0 \rightarrow \check{\mathcal{T}} \xrightarrow{\eta} \text{Pic}(\mathcal{E}) \xrightarrow{\phi} E(\widehat{K}) \rightarrow 0 \quad (9)$$

is a short exact sequence of Abelian groups.

Proof.

1. Notice that the morphism ϕ is surjective: given any $\widehat{K}$ -rational divisor C on E , taking the schematic closure of its irreducible components gives a divisor $\bar{C}$ on $\mathcal{E}$ such that $\bar{C}.E = C$.

2. By Remark (4.1), $\check{\mathcal{T}} \subset \ker(\phi)$. To show that $\check{\mathcal{T}} = \ker(\phi)$, consider $\Upsilon \in \ker(\phi)$, i.e., $\Upsilon = cl(D)$, where $D|_E \sim 0$ on E . But then $D|_E = \text{div}(h)$, where

$$h \in \widehat{K}(E) = \bar{k}(\mathcal{S})(E) = \bar{k}(\mathcal{E}),$$

and hence there exists $H \in \bar{k}(\mathcal{E})$ such that $(H)|_E = (h)$.

If $D' := D - (H)$, then D' must be in some fiber, i.e., $D' \in \check{\mathcal{T}}$, and therefore

$$\Upsilon = cl(D) = cl(D') \in \check{\mathcal{T}}.$$

♠

Theorem 4.2 (A Shioda-Tate Formula for Elliptic N -folds) *Embed $\mathcal{E}(\mathcal{S}/\bar{k})$ into $\text{NS}(\mathcal{E})$ by sending a section σ to the divisor $\overline{\sigma(\mathcal{S})} - \sigma_0(\mathcal{S})$ (where $\overline{\sigma(\mathcal{S})}$ is the schematic closure of $\sigma(\mathcal{S})$ in $\mathcal{E}$).*

Then there is a decomposition of $\text{Gal}(\bar{k}/k)$ -modules,

$$\text{NS}(\mathcal{E}) \otimes \mathbb{Q} \cong (\mathcal{E}(\mathcal{S}/\bar{k}) \otimes \mathbb{Q}) \oplus \mathcal{T}.$$

Proof. Comparing the short exact sequences (3) and (9), we see that ψ maps $\text{Pic}(\mathcal{E})$ surjectively onto $E(\widehat{K})$, while at the same time sending $\text{Pic}^0(\mathcal{E})$ to $B(\bar{k}) = 0$. This implies that

$$\text{NS}(\mathcal{E}) := \text{Pic}(\mathcal{E})/\text{Pic}^0(\mathcal{E}) \twoheadrightarrow E(\widehat{K}), \quad (10)$$

with kernel $\mathcal{T}'$, the image of $\check{\mathcal{T}}$ in $\text{NS}(\mathcal{E})$. Thus, we have an exact sequence

$$0 \longrightarrow \mathcal{T}' \longrightarrow \text{NS}(\mathcal{E}) \longrightarrow E(\widehat{K}) \longrightarrow 0.$$

Since the action of Galois sends vertical divisors to vertical divisors, and horizontal to horizontal, this sequence splits as a Galois module after tensoring with $\mathbb{Q}$; noting that $\mathcal{T}' \otimes \mathbb{Q} = \mathcal{T}$ then gives the desired formula. ♠

Corollary 4.1

$$\text{rank NS}(\mathcal{E}/k) = 1 + \text{rank } \mathcal{E}(\mathcal{S}/k) + \text{rank NS}(\mathcal{S}/k) + \text{rank } \mathcal{F}^{\text{Gal}(\bar{k}/k)},$$

where $\mathcal{F}$ is the vector space generated by the non-identity geometrically irreducible components of the fibral divisors.

Proof. Take $\text{Gal}(\bar{k}/k)$ -invariants of the Shioda-Tate formula for elliptic n -folds. This gives

$$\text{rank NS}(\mathcal{E}/k) = \text{rank } \mathcal{E}(\mathcal{S}/k) + \text{rank } \mathcal{T}^{\text{Gal}(\bar{k}/k)}.$$

But recall (see Definition 2.14 ff) that $\mathcal{T}$ is generated by (0) , $\pi^*(\text{NS}(\mathcal{S}))$, and $\mathcal{F}$. Therefore,

$$\text{rank } \mathcal{T}^{\text{Gal}(\bar{k}/k)} = 1 + \text{rank NS}(\mathcal{S}/k) + \text{rank } \mathcal{F}^{\text{Gal}(\bar{k}/k)}$$

as required. ♠

5 The Singular Fibers

The main goal of this section is to prove Theorem 5.1 below, which establishes a geometric interpretation for the action of Frobenius on the singular fibers. Our main tools will be Tate's Algorithm for determining the singularity-type of a given fiber, and an effective version of the Geometric Tchebotarev Density Theorem, and this requires that we now restrict to the case of an **elliptic threefold** $\mathcal{E}/k$.

Theorem 5.1 *Let $\mathcal{E}/k$ be an elliptic threefold, with notation as before. Then*

$$\sum_{x \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} (m_x - 1) = q_{\mathfrak{p}} \text{Trace}(\text{Frob}_{\mathfrak{p}}|_{\tilde{\mathcal{F}}}) + O(\sqrt{q_{\mathfrak{p}}}),$$

where $\tilde{\mathcal{F}}$ is the vector space generated by all non-identity components of $\pi^{-1}(\tilde{\Delta})$ (the identity component is the component intersecting (0)).

Proof.

1. Recall (Definition 2.5) that if $\Delta = \Delta_1 + \cdots + \Delta_r$ is the irreducible decomposition of the discriminant Δ of $\pi : \mathcal{E} \rightarrow \mathcal{S}$, then, for all $\mathfrak{p} \notin B$, the discriminant locus of $\tilde{\pi} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{S}}$ is given by $\tilde{\Delta}$, the reduction of $\Delta \bmod(\mathfrak{p})$, which has irreducible decomposition

$$\tilde{\Delta} = \tilde{\Delta}_1 + \cdots + \tilde{\Delta}_r.$$

From this it follows that, for all $\mathfrak{p} \notin B$ (where we recall, from section 2.5, that B is a finite set of “bad” primes), genus $\tilde{\Delta}_i = \text{genus } \Delta_i$ for all i .

2. Now assume that, for all $1 \leq j \leq r$:

$$\sum_{x \in \tilde{\Delta}_j(\mathbb{F}_{\mathfrak{p}})} (m_x - 1) = q_{\mathfrak{p}} \text{Trace}(\text{Frob}_{\mathfrak{p}}|_{\tilde{\mathcal{F}}_j}) + O(\sqrt{q_{\mathfrak{p}}}),$$

where $\tilde{\mathcal{F}}_j$ is the vector space generated by the non-identity fibral divisors over $\tilde{\Delta}_j$. Then

$$\begin{aligned} \sum_{x \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} (m_x - 1) &= \sum_{j=1}^r \sum_{x \in \tilde{\Delta}_j(\mathbb{F}_{\mathfrak{p}})} (m_x - 1), \\ &= \sum_{j=1}^r q_{\mathfrak{p}} \text{Trace}(\text{Frob}_{\mathfrak{p}}|_{\tilde{\mathcal{F}}_j}) + O(\sqrt{q_{\mathfrak{p}}}), \\ &= q_{\mathfrak{p}} \text{Trace}(\text{Frob}_{\mathfrak{p}}|_{\tilde{\mathcal{F}}}) + O(\sqrt{q_{\mathfrak{p}}}). \end{aligned}$$

Thus, it suffices to prove the theorem with Δ replaced by one of its irreducible components.

3. The singular points on Δ are determined by the simultaneous vanishing of a finite set of polynomials. But, for a given polynomial $f(D) \in k[\Delta]$, there are only finitely many $x \in \Delta$ such that $f(x) \neq 0$, but $f(x) \equiv 0 \pmod{\mathfrak{p}}$ for all primes $\mathfrak{p}$. Thus, there can only be finitely-many nonsingular $x \in \Delta$ which are singular points on $\tilde{\Delta}$ for all reductions $\text{mod}(\mathfrak{p})$.

Let G be the set containing all singular points on Δ and all points which are singular $\text{mod}(\mathfrak{p})$ for all $\mathfrak{p}$.

At the singular points of $s \in \tilde{\Delta}$, we note that the number of components lying over $\tilde{\mathcal{E}}_s$ is bounded by the number of components lying over $\mathcal{E}_s$ (this follows from Section 2.5, Statement 4), hence is bounded by a constant independent of $\mathfrak{p}$. Furthermore, the following argument will show that the number of singular points on $\tilde{\Delta}$ is bounded independently of $\mathfrak{p}$, and therefore singular points and the fibers over them contribute only to the error term in Theorem 5.1.

Denote by $\eta: \hat{\Delta} \rightarrow \Delta$ the desingularization of Δ . If Δ_{ns} is the set of nonsingular points on Δ , then

$$\eta^{-1}(\Delta_{ns}) \cong \Delta_{ns},$$

while, for the finitely-many points $\{s_g\}_1^R \in G$, $\#\eta^{-1}(s_g) = S_g$, a finite constant.

For almost all $\mathfrak{p}$, $\hat{\Delta} = \tilde{\Delta}$, and $\#\eta^{-1}(s_g) = \#\eta^{-1}(\tilde{s}_g)$. Thus, letting $S := \max_g \{S_g\}$, we see that for all primes $\mathfrak{p} \notin B$,

$$|\#\hat{\Delta}(\mathbb{F}_{\mathfrak{p}}) - \#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})| \leq R \cdot S.$$

4. The next step is to show that at almost every point $x \in \tilde{\Delta}$, the singular fiber $\pi^{-1}(x)$ is of Kodaira type, and that the singularity type is locally constant. This can be accomplished by considering the localization of $\tilde{\mathcal{E}}$ at $\tilde{\Delta}$, and applying Tate's Algorithm.

Let $\mathcal{O}_{\tilde{\mathcal{S}}}$ be the structure sheaf of $\tilde{\mathcal{S}}$, and $\mathcal{O}_{\tilde{\mathcal{S}}, \tilde{\Delta}}$ the local ring of $\tilde{\Delta}$ on $\tilde{\mathcal{S}}$. Then $R := \mathcal{O}_{\tilde{\mathcal{S}}, \tilde{\Delta}}$ is a DVR, with residue field $F := \mathbb{F}_{\mathfrak{p}}(\tilde{\Delta}) = \mathbb{F}_{\mathfrak{p}}(D)$, and prime $\mathfrak{m}$; let

$$\tilde{\mathcal{E}}_{\tilde{\Delta}} := \tilde{\mathcal{E}} \times_{\mathcal{O}_{\tilde{\mathcal{S}}}} \mathcal{O}_{\tilde{\mathcal{S}}, \tilde{\Delta}}$$

be the localization of $\tilde{\mathcal{E}}$ at $\tilde{\Delta}$; then a Weierstrass Equation for $\tilde{\mathcal{E}}_{\tilde{\Delta}}$ is given by:

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6, \quad a_i \in R = \mathbb{F}_{\mathfrak{p}}[D]. \quad (11)$$

Remark 5.1 Notice that $\tilde{\mathcal{E}}_{\tilde{\Delta}}$ is an elliptic curve defined over a DVR R with non-perfect residue field F , whereas Tate's Algorithm is for elliptic curves over DVR's with perfect residue fields. However, the proof works verbatim for any residue field with the property that all extensions of degree 2 or 3 are separable. (For the proof of Tate's Algorithm, see Tate's original paper [31], or Silverman [29] IV.9). Thus, if B is expanded to include all primes $\mathfrak{p}$ such that $2|q_{\mathfrak{p}}$ or $3|q_{\mathfrak{p}}$, then Tate's Algorithm can also be applied here.

5. Localizing $\mathcal{E}$ over Δ gives an elliptic curve defined over the DVR $\mathcal{O}_{S,\Delta}$, with perfect residue field $k(\Delta)$ - hence a straightforward application of Tate's Algorithm shows that the Kodaira singularity type of $\mathcal{E}$ is locally constant over Δ .

Then, for all but finitely-many $\mathfrak{p}$,

- (a) The Kodaira type of $\mathcal{E}$ over Δ is the same as the Kodaira type of $\tilde{\mathcal{E}}$ over Δ , call it $\mathfrak{D}$.

- (b) Let

$$\Delta_{KF} := \{x \in \Delta \mid x \notin G \text{ and } \mathcal{E}_x \text{ has Kodaira type } \mathfrak{D}\}.$$

$$\text{Then } \Delta_{KF} = \tilde{\Delta}_{KF}.$$

The first statement follows because Tate's Algorithm is applied to the local equations of $\mathcal{E}$ and $\tilde{\mathcal{E}}$ over Δ and $\tilde{\Delta}$ respectively, and for almost all $\mathfrak{p}$, they will be the same equation, with the same zeros.

The second statement follows from Elimination Theory arguments.

6. The proof now reduces to an examination of each fiber type.

Note first that, since we are considering the action of $\text{Frob}_{\mathfrak{p}}$ on a subspace of $\text{NS}(\tilde{\mathcal{E}}/\bar{\mathbb{F}}_{\mathfrak{p}})$, we must look at all fibral components that are irreducible over $\bar{\mathbb{F}}_{\mathfrak{p}}$ (and not over $\bar{F}$!). $\text{Trace}(\text{Frob}_{\mathfrak{p}})$ picks out from among these the ones that are defined over $\mathbb{F}_{\mathfrak{p}}$. These will be called $\mathbb{F}_{\mathfrak{p}}$ -rational components.

Following Tate's notation, we now define

$$a_{i,j} := \mathfrak{m}^{-j} a_i.$$

Consider the divisor $\tilde{\pi}^{-1}(\tilde{\Delta})$. Let

$$\begin{aligned} H &= \text{number of irreducible components over } \bar{F}, \\ h &= \text{number of irreducible components over } \bar{F} \text{ that are defined over } \mathbb{F}_{\mathfrak{p}}, \\ M &= \text{number of } \mathbb{F}_{\mathfrak{p}}\text{-rational components.} \end{aligned}$$

- Type I_n , $n > 0$.

There are n components of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$, so $H = n$. According to Tate's Algorithm, the number M of $\mathbb{F}_{\mathfrak{p}}$ -rational components is determined by the splitting field F' of

$$P(T) = T^2 + a_1 T - a_2.$$

- (a) If $F' = F$, then $M = n$; furthermore, $F' = F$ implies that, for all $d \in \tilde{\Delta}$,

$$P_d(T) = T^2 + a_1(d)T - a_2(d)$$

splits in $\mathbb{F}_{\mathfrak{p}}$. Therefore we also have $m_d = n$ for every $d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})$.

- (b) If $F' \neq F$ and $F' = L(\tilde{\Delta})$ for L some finite extension of $\mathbb{F}_p$, then in fact the n components of $\pi^{-1}(\tilde{\Delta})$ are defined over $\bar{\mathbb{F}}_p$, but they are not all defined over $\mathbb{F}_p$ because $F' \neq F$. To determine the number of $\mathbb{F}_p$ -rational components, we consider what happens at the fibers over points $d \in \tilde{\Delta}(\mathbb{F}_p)$.

$F' = L(\tilde{\Delta})$ implies that, for all $d \in \tilde{\Delta}(\mathbb{F}_p)$, the splitting field of $P_d(T)$ will be $L \neq \mathbb{F}_p$. Hence, the Galois action reflects the geometric components of the polygon. This means there will always be two invariant components when n is even, and one invariant component when n is odd. Therefore we have:

for even n : $M = m_d = 2$, for all $d \in \tilde{\Delta}(\mathbb{F}_p)$,

for odd n : $M = m_d = 1$, for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- (c) Otherwise, from the discussion in part (b) it is clear that $h = 2$ for even n , $h = 1$ for odd n , and

$$\begin{aligned} M &= \frac{n+2}{2} = \frac{H+h}{2} \quad \text{for even } n, \\ M &= \frac{n+1}{2} = \frac{H+h}{2} \quad \text{for odd } n. \end{aligned}$$

This can be explained as follows: since $F' \neq L(\tilde{\Delta})$, the irreducible components of $\pi^{-1}(\tilde{\Delta})$ are NOT defined over $\bar{\mathbb{F}}_p$. Thus, the $\bar{\mathbb{F}}_p$ -components are exactly the $\mathbb{F}_p$ -components, and consist of the Frob_p orbits. Since the action on I_n is reflection, there will be $\frac{n+2}{2}$ for even n , and $\frac{n+1}{2}$ for odd n .

At the level of fibers over points $d \in \tilde{\Delta}(\mathbb{F}_p)$, m_d alternates between 1 and n (in the case n odd), and between 2 and n (if n even), depending on whether or not $\mathbb{F}_p$ is the splitting field of $P_d(T)$.

- Type II. There is only one component of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$; since this includes the identity component Θ_0 , which is defined over F , we get $M = m_d = 1$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.
- Type III. There are two components of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$. One is Θ_0 , so the other must also remain fixed under the action of Frobenius. Hence $M = m_d = 2$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.
- Type IV. There are three components of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$, so $H = 3$. Let F' be the splitting field of

$$P(T) = T^2 + a_{3,1}T - a_{6,2}.$$

- (a) If $F' = F$, then $M = m_d = 3$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.
- (b) If $F' \neq F$ and $F' = L(\tilde{\Delta})$, where L is a finite extension of $\mathbb{F}_p$, then $M = m_d = 1$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.
- (c) Otherwise, m_d alternates between 1 and 3, which shows that $h = 1$, and $M = 2 = \frac{H+h}{2}$.

- Type I_n^* .

There are $n + 5$ components of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$. To determine the number of $\mathbb{F}_p$ -rational components, consider the polynomial

$$Q(T) = T^3 + a_{2,1}T^2 + a_{4,2}T + a_{6,3}. \quad (12)$$

- (a) If $Q(T)$ has three distinct roots in $\bar{F}$, then we are in the case I_0^* . There are 5 components defined over $\bar{F}$ (i.e., $H = 5$). Let F' be the splitting field of $Q(T)$.

- i. If $F' = F$, then $M = m_d = 5$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- ii. If $Q(T)$ is reducible in F , but $F' \neq F$, then let $P_1(T)$ be the irreducible part of $Q(T)$, and let F'' be the splitting field of $P_1(T)$.

If $F'' = L(\tilde{\Delta})$, where L is a finite extension of $\mathbb{F}_p$, then $M = M_d = 3$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

If $F'' \neq L(\tilde{\Delta})$, then $M = 4$, and m_d alternates between 3 and 5.

- iii. If $Q(T)$ is irreducible in F , and $F' = L(\tilde{\Delta})$, where L is a finite extension of $\mathbb{F}_p$, then $M = m_d = 2$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- iv. If $Q(T)$ is irreducible in F , and $F' \neq L(\tilde{\Delta})$, then m_d alternates between 2, 3, and 5, (so $h = 2$), and $M = 3 = \frac{H+h}{2}$.

- (b) If $Q(T)$ has one simple root and one double root in $\bar{F}$, then we have type I_n^* , $n \geq 1$. In this case, $H = n + 5$, and the number of $\mathbb{F}_p$ -rational components ($n + 3$ or $n + 5$) is determined by the splitting of a quadratic polynomial $P(T)$. Let C be its splitting field.

- i. If $C = F$, then $M = m_d = n + 5$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- ii. If $C = L(\tilde{\Delta})$, where L is a finite extension of $\mathbb{F}_p$, then $M = m_d = n + 3$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- iii. If $C \neq L(\tilde{\Delta})$, then m_d alternates between $n + 3$, and $n + 5$ (so $h = n + 3$), and $M = n + 4 = \frac{H+h}{2}$.

- Type III^* .

There are eight components of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$, and they are all fixed by the action of Frobenius. Hence, $M = m_d = 8$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- Type II^* .

There are nine components of $\tilde{\pi}^{-1}(\tilde{\Delta})$ defined over $\bar{F}$, and they are all fixed by the action of Frobenius. Hence, $M = m_d = 9$ for all $d \in \tilde{\Delta}(\mathbb{F}_p)$.

- Type IV^* .

There are seven components defined over $\bar{F}$, therefore $H = 7$.

Let F' be the splitting field of

$$P(T) = T^2 + a_{3,2}T - a_{6,4}.$$

- (a) If $F' = F$ then $M = m_d = 7$ for all $d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})$.
- (b) If $F' = L(\tilde{\Delta})$, where L is a finite extension of $\mathbb{F}_{\mathfrak{p}}$, then $M = m_d = 3$ for all $d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})$.
- (c) If $F' \neq L(\tilde{\Delta})$, then m_d alternates between 3 and 7 (so $h = 3$), while $M = 5 = \frac{H+h}{2}$.

7. The proof of the theorem now reduces to analyzing two cases:

- (a) The case $M = m_d$ for all $d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})$:

We have

$$\begin{aligned}
\sum_{d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} (m_d - 1) &= \sum_{d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} (M - 1) \\
&= \#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})(M - 1) \\
&= \#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}})
\end{aligned} \tag{13}$$

- (b) The case $M \neq m_d$: This requires a Chebotarev argument.

We use the following effective version of the geometric Chebotarev Density Theorem:

Theorem 5.2 (Murty, Scherk [21]) *If $X \rightarrow Y$ is a geometric covering of curves over $\mathbb{F}_{\mathfrak{p}}$ (i.e., Y is defined over $\mathbb{F}_{\mathfrak{p}}$, and $\mathbb{F}_{\mathfrak{p}}$ is the algebraic closure of itself in $\mathbb{F}_{\mathfrak{p}}(X)$), then*

$$\left| \psi_C(r) - \frac{|C|}{|G|} \psi(r) \right| \leq 2g_X \frac{|C|}{|G|} q_{\mathfrak{p}}^{\frac{r}{2}} + |D|; \tag{14}$$

where, for $y \in Y$ unramified, C is a conjugacy class in $G := \text{Gal}(X/Y)$, σ_y is a Frobenius conjugacy class of y , and

$$\begin{aligned}
Y_r &= Y \times_{\mathbb{F}_{\mathfrak{p}}} \mathbb{F}_{\mathfrak{p}^r}, \\
\bar{Y} &= Y \times_{\mathbb{F}_{\mathfrak{p}}} \bar{\mathbb{F}}_{\mathfrak{p}}, \\
\psi_C(r) &= \#\{y \in Y_r | y \text{ unramified}, \sigma_y \in C\}, \\
\psi(r) &= \#\{y \in Y_r | y \text{ unramified}\}, \\
D &= \text{the set of ramified points in } \bar{Y}.
\end{aligned}$$

- (c) To apply this version of Chebotarev, we note that in our case, X is the curve defined by the polynomial $P(T)$, $Y = \tilde{\Delta}$, and, since we are considering only points in $\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})$, $r = 1$. Thus, we get the inequality:

$$\begin{aligned}
|\psi_C(1) - \frac{|C|}{|G|} \psi(1)| &\leq 2g_X \frac{|C|}{|G|} \sqrt{q_{\mathfrak{p}}} + |D|, \\
&\leq 2g_X \sqrt{q_{\mathfrak{p}}} + |D|.
\end{aligned}$$

Since, furthermore, by the Riemann-Hurwitz formula

$$2g_X - 2 = (2g_Y - 2)|G| + |D|,$$

the inequality becomes

$$\begin{aligned} |\psi_C(1) - \frac{|C|}{|G|}\psi(1)| &\leq 2g_Y\sqrt{q_{\mathfrak{p}}} + (2 - 2|G| + |D|)\sqrt{q_{\mathfrak{p}}} + |D|; \\ &\leq K\sqrt{q_{\mathfrak{p}}} + |D|. \end{aligned}$$

where K is a constant depending only on $|G|$, g_Y and $|D|$, and hence is bounded independently of $\mathfrak{p}$. This gives

$$\psi_C(1) = \frac{|C|}{|G|}\psi(1) + O(\sqrt{q_{\mathfrak{p}}}). \quad (15)$$

Noting that $\psi(1) = \#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}}) - |D|$, gives

$$\psi_C(1) = \frac{|C|}{|G|}\#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}}) + O(\sqrt{q_{\mathfrak{p}}}).$$

Next, we want to observe what happens for $\deg(P) = 2$, and $\deg(P) = 3$.

i. $\deg(P) = 2$.

In this case, $|G| = 2$; one of the two conjugacy classes consists of id , and the other is a transposition t .

If, for a given $d \in \tilde{\Delta}$, $\sigma_d = id$, then $m_d = H$. If $\sigma_d = t$, then $m_d = h$. Thus, the sum of components of fibers breaks up into:

$$\begin{aligned} \sum_{d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} m_d &= \sum_{d: \sigma_d = id} m_d + \sum_{d: \sigma_d = t} m_d \\ &= \sum_{d: \sigma_d = id} H + \sum_{d: \sigma_d = t} h \\ &= H|\psi_{id}| + h|\psi_t| \\ &= H\frac{1}{2}\#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}}) + h\frac{1}{2}\#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}}) + O(\sqrt{q_{\mathfrak{p}}}) \\ &= \frac{H+h}{2}\#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}}) + O(\sqrt{q_{\mathfrak{p}}}). \end{aligned}$$

ii. $\deg(P) = 3$.

In this case, $|G| = 3$ or $|G| = 6$.

A. If $|G| = 3$, there are three conjugacy classes corresponding to the three elements of G : id , v_1 , and v_2 .

If, for a given $d \in \tilde{\Delta}$, $\sigma_d = id$, then $m_d = 5$; if $\sigma_d = v_i$, $i = 1, 2$, then $m_d = 2$. Thus, the sum of components of fibers breaks up into:

$$\begin{aligned}
\sum_{d \in \tilde{\Delta}(\mathbb{F}_p)} m_d &= \sum_{d: \sigma_d = id} m_d + \sum_{d: \sigma_d = v_1} m_d + \sum_{d: \sigma_d = v_2} m_d \\
&= \sum_{d: \sigma_d = id} 5 + \sum_{d: \sigma_d = v_1} 2 + \sum_{d: \sigma_d = v_2} 2 \\
&= 5|\psi_{id}| + 2|\psi_{v_1}| + 2|\psi_{v_2}| \\
&= 5\left(\frac{1}{3}\right)(\#\tilde{\Delta}(\mathbb{F}_p)) + 2\left(\frac{1}{3}\right)(\#\tilde{\Delta}(\mathbb{F}_p)) \\
&\quad + 2\left(\frac{1}{3}\right)(\#\tilde{\Delta}(\mathbb{F}_p)) + O(\sqrt{q_p}) \\
&= 3\#\tilde{\Delta}(\mathbb{F}_p) + O(\sqrt{q_p}).
\end{aligned}$$

B. If $|G| = 6$, then there are three conjugacy classes; one conjugacy class consists of id ; a second one is a transposition t , and this has 3 elements; the third conjugacy class is the even permutations v , of which there are two.

If, for a given $d \in \tilde{\Delta}$, $\sigma_d = id$, then $m_d = 5$. If $\sigma_d = t$, then $m_d = 3$, and if $\sigma_d = v$, then $m_d = 2$. Thus, the sum of components of fibers breaks up into:

$$\begin{aligned}
\sum_{d \in \tilde{\Delta}(\mathbb{F}_p)} m_d &= \sum_{d: \sigma_d = id} m_d + \sum_{d: \sigma_d = t} m_d + \sum_{d: \sigma_d = v} m_d \\
&= \sum_{d: \sigma_d = id} 5 + \sum_{d: \sigma_d = t} 3 + \sum_{d: \sigma_d = v} 2 \\
&= 5|\psi_{id}| + 3|\psi_t| + 2|\psi_v| \\
&= 5\left(\frac{1}{6}\right)(\#\tilde{\Delta}(\mathbb{F}_p)) + 3\left(\frac{3}{6}\right)(\#\tilde{\Delta}(\mathbb{F}_p)) \\
&\quad + 2\left(\frac{2}{6}\right)(\#\tilde{\Delta}(\mathbb{F}_p)) + O(\sqrt{q_p}) \\
&= 3\#\tilde{\Delta}(\mathbb{F}_p) + O(\sqrt{q_p}).
\end{aligned}$$

(d) By Weil's Estimate, we have:

$$\#\tilde{\Delta}(\mathbb{F}_p) = 1 + a_p(\Delta) + q_p,$$

where, if g_Δ is the genus of Δ ,

$$|a_p(\Delta)| \leq \sqrt{q_p} g_\Delta.$$

Furthermore, $\text{Trace}(\text{Frob}_p | \tilde{\mathcal{F}}) < M$ is bounded independently of p , because for almost all p , M is given by the number of irreducible components of the singular

fiber over Δ . Therefore,

$$\begin{aligned}
\#\tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}}) &= (1 + a_{\mathfrak{p}}(\Delta) + q_{\mathfrak{p}})\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}}) \\
&= q_{\mathfrak{p}}\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}}) \\
&\quad + (1 + a_{\mathfrak{p}}(\Delta))\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}}) \\
&= q_{\mathfrak{p}}\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}}) + O(\sqrt{q_{\mathfrak{p}}}).
\end{aligned}$$

Thus, combining this with the result in equation (13) gives

$$\sum_{d \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} (m_d - 1) = q_{\mathfrak{p}}\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}}) + O(\sqrt{q_{\mathfrak{p}}}).$$

♠

6 The Main Theorem

We now have all the information necessary to prove Theorem 1.1. What remains is to run it through the L -series machinery, and apply Tate's Conjecture.

Recall that, for a smooth variety $\mathcal{V}/k$, the **Hasse-Weil L-series** attached to $H_{\text{ét}}^i(\mathcal{V}/\bar{k})$, denoted $L_2(\mathcal{V}, s)$, is given by

$$L_2(\mathcal{V}, s) := \prod_{\mathfrak{p}} P_{\mathfrak{p},2}(\tilde{\mathcal{V}}, q_{\mathfrak{p}}^{-s}).$$

where

$$P_{\mathfrak{p},i} = \det \left(1 - \text{Frob}_{\mathfrak{p}} t | H_{\text{ét}}^i(\mathcal{V}/\bar{k}; \mathbb{Q}_l) \right).$$

Remark 6.1 *To be precise, since in this paper we are working over all primes $\mathfrak{p} \in U$ (i.e., we are excluding the primes in B),*

$$\begin{aligned}
L_2(\mathcal{E}, s) &\approx \prod_{\mathfrak{p}} P_{\mathfrak{p},2}(\tilde{\mathcal{E}}, q_{\mathfrak{p}}^{-s}) \quad \text{and} \\
L_2(\mathcal{S}, s) &\approx \prod_{\mathfrak{p}} P_{\mathfrak{p},2}(\tilde{\mathcal{S}}, q_{\mathfrak{p}}^{-s}),
\end{aligned}$$

where the symbol $\approx$ is used to indicate that the two sides agree up to finitely many Euler factors. This, however, has no effect on the residue computation.

Conjecture 6.1 (Tate's Conjecture) ([30], Conjecture 2). *Let $\mathcal{V}$ be a smooth projective variety defined over k , and let $L_2(\mathcal{V}, s)$ be the Hasse-Weil L -function attached to $H_{\text{ét}}^2(\mathcal{V}/\bar{k})$. Then $L_2(\mathcal{V}, s)$ has a meromorphic continuation to $\mathbb{C}$, and has a pole at $s = 2$ of order:*

$$-\text{ord}_{s=2} L_2(\mathcal{V}, s) = \text{rank NS}(\mathcal{E}/k).$$

Finally, we are ready to prove the main theorem:

Theorem 1.1 *Let $\mathcal{E} \rightarrow \mathcal{S}$ be a non-split elliptic threefold defined over a number field k . Then Tate's Conjecture for $\mathcal{E}/k$ and $\mathcal{S}/k$ implies*

$$\operatorname{res}_{s=1} \sum_{\mathfrak{p}} -A_{\mathfrak{p}}(\mathcal{E}) \frac{\log \mathfrak{p}}{\mathfrak{p}^s} = \operatorname{rank} \mathcal{E}(\mathcal{S}/k).$$

Proof.

1. Look at $\tilde{\mathcal{E}}$ as a fibration of curves, and use the Lefschetz Fixed-Point Theorem to count its rational points fiber by fiber:

$$\begin{aligned}
\#\tilde{\mathcal{E}}(\mathbb{F}_{\mathfrak{p}}) &= \sum_{x \in \tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}})} \#\tilde{\mathcal{E}}_x(\mathbb{F}_{\mathfrak{p}}) \\
&= \sum_{x \in \tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}})} (1 - a_{\mathfrak{p}}(\tilde{\mathcal{E}}_x) + q_{\mathfrak{p}} + (m_x - 1)q_{\mathfrak{p}}) \\
&= (1 + q_{\mathfrak{p}})\#\tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}}) - q_{\mathfrak{p}}^2 A_{\mathfrak{p}}(\mathcal{E}) + \sum_{x \in \tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}})} (m_x - 1)q_{\mathfrak{p}}. \tag{16}
\end{aligned}$$

Since $m_x = 1$ for all non-singular fibers $\tilde{\mathcal{E}}_x$ with $x \in \tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}})$,

$$\sum_{x \in \tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}})} (m_x - 1)q_{\mathfrak{p}} = \sum_{x \in \tilde{\Delta}(\mathbb{F}_{\mathfrak{p}})} (m_x - 1)q_{\mathfrak{p}}.$$

Therefore, combining equation (16) with the result of Theorem 5.1 gives

$$\begin{aligned}
\#\tilde{\mathcal{E}}(\mathbb{F}_{\mathfrak{p}}) &= (1 + q_{\mathfrak{p}})\#\tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}}) - q_{\mathfrak{p}}^2 A_{\mathfrak{p}}(\mathcal{E}) \\
&\quad + q_{\mathfrak{p}} \text{Trace}(\text{Frob}_{\mathfrak{p}}|_{\tilde{\mathcal{F}}})q_{\mathfrak{p}} + O(\sqrt{q_{\mathfrak{p}}^3}). \tag{17}
\end{aligned}$$

2. Now use the Lefschetz Fixed-Point Theorem again, this time considering $\mathcal{E}$ as a three-fold, and $\mathcal{S}$ as a surface:

$$\#\tilde{\mathcal{E}}(\mathbb{F}_{\mathfrak{p}}) = 1 - a_{\mathfrak{p}}(\tilde{\mathcal{E}}) + b_{\mathfrak{p}}(\tilde{\mathcal{E}}) - c_{\mathfrak{p}}(\tilde{\mathcal{E}}) + q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{E}}) - q_{\mathfrak{p}}^2 a_{\mathfrak{p}}(\tilde{\mathcal{E}}) + q_{\mathfrak{p}}^3. \tag{18}$$

$$\#\tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}}) = 1 - a_{\mathfrak{p}}(\tilde{\mathcal{S}}) + b_{\mathfrak{p}}(\tilde{\mathcal{S}}) - q_{\mathfrak{p}}a_{\mathfrak{p}}(\tilde{\mathcal{S}}) + q_{\mathfrak{p}}^2, \tag{19}$$

where we note that

$$\begin{aligned}
\text{Trace}(\text{Frob}_{\mathfrak{p}}|H_{\text{ét}}^5(\mathcal{E}, \mathbb{Q}_l)) &= q_{\mathfrak{p}}^2 a_{\mathfrak{p}}(\tilde{\mathcal{E}}), \\
\text{Trace}(\text{Frob}_{\mathfrak{p}}|H_{\text{ét}}^4(\mathcal{E}, \mathbb{Q}_l)) &= q_{\mathfrak{p}} b_{\mathfrak{p}}(\tilde{\mathcal{E}}), \\
\text{Trace}(\text{Frob}_{\mathfrak{p}}|H_{\text{ét}}^3(\mathcal{S}, \mathbb{Q}_l)) &= q_{\mathfrak{p}} a_{\mathfrak{p}}(\tilde{\mathcal{S}}),
\end{aligned}$$

are given by duality

$$\begin{aligned}
H_{\text{ét}}^1(\mathcal{E}, \mathbb{Q}_l) &\cong H_{\text{ét}}^5(\mathcal{E}, \mathbb{Q}_l)^{\vee}, \\
H_{\text{ét}}^2(\mathcal{E}, \mathbb{Q}_l) &\cong H_{\text{ét}}^4(\mathcal{E}, \mathbb{Q}_l)^{\vee}, \\
H_{\text{ét}}^1(\mathcal{S}, \mathbb{Q}_l) &\cong H_{\text{ét}}^3(\mathcal{S}, \mathbb{Q}_l)^{\vee}.
\end{aligned}$$

Since $a_{\mathfrak{p}}(\mathcal{S}) = a_{\mathfrak{p}}(\mathcal{E})$ by Corollary (3.1), equation (18) implies:

$$\#\tilde{\mathcal{E}}(\mathbb{F}_{\mathfrak{p}}) = 1 - a_{\mathfrak{p}}(\tilde{\mathcal{S}}) + b_{\mathfrak{p}}(\tilde{\mathcal{E}}) - c_{\mathfrak{p}}(\tilde{\mathcal{E}}) + q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{E}}) - q_{\mathfrak{p}}^2 a_{\mathfrak{p}}(\tilde{\mathcal{S}}) + q_{\mathfrak{p}}^3. \tag{20}$$

A different expression for the number of rational points on $\tilde{\mathcal{E}}$ is obtained by combining equations (17) and (19):

$$\begin{aligned} \#\tilde{\mathcal{E}}(\mathbb{F}_{\mathfrak{p}}) &= 1 + q_{\mathfrak{p}} + q_{\mathfrak{p}}^2 + q_{\mathfrak{p}}^3 - a_{\mathfrak{p}}(\tilde{\mathcal{S}}) - 2q_{\mathfrak{p}}a_{\mathfrak{p}}(\tilde{\mathcal{S}}) - q_{\mathfrak{p}}^2a_{\mathfrak{p}}(\tilde{\mathcal{S}}) + b_{\mathfrak{p}}(\tilde{\mathcal{S}}) + q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{S}}) \\ &\quad - q_{\mathfrak{p}}^2A_{\mathfrak{p}}(\mathcal{E}) + q_{\mathfrak{p}}\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}})q_{\mathfrak{p}} + O(\sqrt{q_{\mathfrak{p}}^3}). \end{aligned} \quad (21)$$

3. Finally, equating the two expressions for the number of rational points on $\tilde{\mathcal{E}}$ in equations (20) and (21) gives an expression for $A_{\mathfrak{p}}(\mathcal{E})$:

$$\begin{aligned} q_{\mathfrak{p}}^2A_{\mathfrak{p}}(\mathcal{E}) &= q_{\mathfrak{p}} - 2q_{\mathfrak{p}}a_{\mathfrak{p}}(\tilde{\mathcal{S}}) + b_{\mathfrak{p}}(\tilde{\mathcal{S}}) + q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{S}}) + c_{\mathfrak{p}}(\tilde{\mathcal{E}}) - b_{\mathfrak{p}}(\tilde{\mathcal{E}}) - q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{E}}) \\ &\quad + q_{\mathfrak{p}}^2 + q_{\mathfrak{p}}\text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}})q_{\mathfrak{p}} + O(\sqrt{q_{\mathfrak{p}}^3}). \end{aligned} \quad (22)$$

By Deligne's Theorem [2], we know, for every smooth, projective variety $\mathcal{V}$ defined over k , that

$$\begin{aligned} |a_{\mathfrak{p}}(\mathcal{V})| &\leq B_1(\mathcal{V})\sqrt{q_{\mathfrak{p}}}, \\ |b_{\mathfrak{p}}(\mathcal{V})| &\leq B_2(\mathcal{V})q_{\mathfrak{p}}, \\ |c_{\mathfrak{p}}(\mathcal{V})| &\leq B_3(\mathcal{V})\sqrt{q_{\mathfrak{p}}^3}, \end{aligned}$$

where $B_i(\mathcal{V}) := \dim H_{\text{ét}}^i(\mathcal{V}/\bar{k}, \mathbb{Q}_l)$ is independent of $\mathfrak{p}$.

Thus, we can group all terms of order $\sqrt{q_{\mathfrak{p}}^3}$ or less together, and obtain:

$$q_{\mathfrak{p}}^2A_{\mathfrak{p}}(\mathcal{E}) = q_{\mathfrak{p}}^2 + q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{S}}) - q_{\mathfrak{p}}b_{\mathfrak{p}}(\tilde{\mathcal{E}}) + \text{Trace}(\text{Frob}_{\mathfrak{p}}|\tilde{\mathcal{F}})q_{\mathfrak{p}}^2 + O(\sqrt{q_{\mathfrak{p}}^3}). \quad (23)$$

4. And now compute residues.

(a)

$$\text{res}_{s=1} \sum_{\mathfrak{p}} \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} = 1. \quad (24)$$

- (b) Letting $L(\mathcal{F}, s)$ be the Artin L -series attached to $\mathcal{F}$, we have, for $\text{Re}(s) > \frac{1}{2}$

$$\begin{aligned} \frac{d}{ds} \log L(\mathcal{F}, s) &= \frac{d}{ds} \sum_{\mathfrak{p}} -\log \det(1 - \text{Frob}_{\mathfrak{p}}q_{\mathfrak{p}}^{-s}|\mathcal{F}) \\ &= \sum_{\mathfrak{p}} -\text{Trace}(\text{Frob}_{\mathfrak{p}}|\mathcal{F}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} + O(1). \end{aligned}$$

Therefore,

$$\begin{aligned}
\operatorname{res}_{s=1} \sum_{\mathfrak{p}} \operatorname{Trace}(\operatorname{Frob}_{\mathfrak{p}} | \tilde{\mathcal{F}}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} &= - \operatorname{res}_{s=1} \frac{d}{ds} \log L(\mathcal{F}, s) \\
&= - \operatorname{ord}_{s=1} L(\mathcal{F}, s) \\
&= \operatorname{rank}(\mathcal{F}^{\operatorname{Gal}(\bar{k}/k)}). \tag{25}
\end{aligned}$$

This last equality follows from [25], Proposition 1.5.1.

(c) Furthermore, for $\operatorname{Re}(s) > \frac{3}{2}$,

$$\begin{aligned}
\frac{d}{ds} \log L_2(\mathcal{E}, s) &= \frac{d}{ds} \sum_{\mathfrak{p}} -\log \det (1 - \operatorname{Frob}_{\mathfrak{p}} q_{\mathfrak{p}}^{-s} | H_{\text{ét}}^2(\mathcal{E}/\bar{k}, \mathbb{Q}_l)) \\
&= \sum_{\mathfrak{p}} -b_{\mathfrak{p}}(\mathcal{E}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} + O(1).
\end{aligned}$$

Therefore,

$$\begin{aligned}
\operatorname{res}_{s=2} \sum_{\mathfrak{p}} b_{\mathfrak{p}}(\mathcal{E}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} &= - \operatorname{res}_{s=1} \frac{d}{ds} \log L_2(\mathcal{E}, s) \\
&= - \operatorname{ord}_{s=2} L_2(\mathcal{E}, s) \\
&= \operatorname{rank} \operatorname{NS}(\mathcal{E}/k) \quad \text{by Tate's Conjecture,} \tag{26}
\end{aligned}$$

and similarly

$$\begin{aligned}
\operatorname{res}_{s=2} \sum_{\mathfrak{p}} b_{\mathfrak{p}}(\mathcal{S}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} &= - \operatorname{res}_{s=1} \frac{d}{ds} \log L_2(\mathcal{S}, s) \\
&= - \operatorname{ord}_{s=2} L_2(\mathcal{S}, s) \\
&= \operatorname{rank} \operatorname{NS}(\mathcal{S}/k) \quad \text{by Tate's Conjecture.} \tag{27}
\end{aligned}$$

5. Combining the residue calculations with equation (23) gives

$$\operatorname{res}_{s=1} \sum_{\mathfrak{p}} -A_{\mathfrak{p}}(\mathcal{E}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} = -1 - \operatorname{rank}(\mathcal{F}^{\operatorname{Gal}(\bar{k}/k)}) - \operatorname{rank} \operatorname{NS}(\mathcal{S}/k) + \operatorname{rank} \operatorname{NS}(\mathcal{E}/k).$$

and by the Shioda-Tate formula for elliptic threefolds (Theorem 4.2), this gives

$$\operatorname{res}_{s=1} \sum_{\mathfrak{p}} -A_{\mathfrak{p}}(\mathcal{E}) \frac{\log q_{\mathfrak{p}}}{q_{\mathfrak{p}}^s} = \operatorname{rank} \mathcal{E}(\mathcal{S}/k). \tag{28}$$

♠

A Notation

$k/\mathbb{Q}$	a number field.
$\bar{k}$	the algebraic closure of k .
$\mathbb{F}_{\mathfrak{p}}$	the residue field of a prime $\mathfrak{p}$ of k .
$\bar{\mathbb{F}}_{\mathfrak{p}}$	the algebraic closure of $\mathbb{F}_{\mathfrak{p}}$.
$q_{\mathfrak{p}}$	the norm of $\mathfrak{p}$, i.e., $q_{\mathfrak{p}} = \#\mathbb{F}_{\mathfrak{p}}$.
$\mathcal{S}/k$	a smooth, projective surface defined over k .
$\mathcal{E}/k$	a (non-split) elliptic threefold $\pi : \mathcal{E} \rightarrow \mathcal{S}$ with section $\sigma : \mathcal{S} \rightarrow \mathcal{E}$; regular, proper and flat over $\mathcal{S}$, and defined over k .
$\tilde{\mathcal{S}}/\mathbb{F}_{\mathfrak{p}}$	the reduction of $\mathcal{S} \pmod{\mathfrak{p}}$.
$\tilde{\mathcal{E}}/\mathbb{F}_{\mathfrak{p}}$	the reduction of $\mathcal{E} \pmod{\mathfrak{p}}$.
K	$= k(\mathcal{S})$, the function field of $\mathcal{S}/k$.
$\hat{K}$	$= \bar{k}(\mathcal{S})$, the function field of $\mathcal{S}/\bar{k}$.
E/K	the generic fiber of $\pi : \mathcal{E} \rightarrow \mathcal{S}$, a smooth elliptic curve defined over K .
(τ, B)	the $\hat{K}/\bar{k}$ -trace of E .
Δ	the discriminant locus of $\mathcal{E} \rightarrow \mathcal{S}$, with irreducible decomposition $\sum_{j=1}^r \Delta_j$.
$\tilde{\Delta}$	the discriminant locus of the fibration $\tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{S}}$ $\pmod{\mathfrak{p}}$, with irreducible decomposition $\sum_{j=1}^r \tilde{\Delta}_j$.
M_j	the number of $\mathbb{F}_{\mathfrak{p}}$ -rational components of $\tilde{\pi}^{-1}(\tilde{\Delta}_j)$.
$\Theta_{i,j}$	$(0 \leq i \leq M_j - 1)$: the $\mathbb{F}_{\mathfrak{p}}$ -rational components of $\tilde{\pi}^{-1}(\tilde{\Delta}_j)$.
$\Theta_{0,j}$	the identity component.
m_x	the number of $\mathbb{F}_{\mathfrak{p}}$ -rational components of the fiber $\tilde{\mathcal{E}}_x$, for given $x \in \tilde{\mathcal{S}}(\mathbb{F}_{\mathfrak{p}})$.

References

- [1] M. Artin. *Théorie des topos et cohomologie étale des schémas*. Lecture notes in mathematics 269; SGA 4. Springer-Verlag, Berlin, 1972.
- [2] P. Deligne. La conjecture de Weil I. *Publ. Math. IHES*, 43:273–307, 1974.
- [3] P. Deligne. La conjecture de Weil II. *Publ. Math. IHES*, 52:313–428, 1981.
- [4] I. Dolgachev and M. Gross. Elliptic threefolds. I. Ogg-Shafarevich theory. *J. Alg. Geom.*, 3:39–80, 1994.
- [5] B. Fisher. Equidistribution Theorems (d’après P. Deligne et N. Katz), Columbia University Number Theory Seminar, New York 1992. *Asterisque*, 228:69–79, 1995.
- [6] E. Freitag and R. Kiehl. *Etale Cohomology and the Weil Conjecture*. Ergebnisse der Mathematik und ihrer Grenzgebiete, 13. Springer Verlag, Berlin, 1987.
- [7] W. Fulton. *Intersection Theory*. Ergebnisse der Mathematik und Ihrer Grenzgebiete, 3. Springer Verlag, 1980.
- [8] W. Fulton. *Introduction to Intersection Theory in Algebraic Geometry*. CBMS Regional Conference Series in Mathematics, no. 54. American Mathematical Society, 1984.
- [9] P. Griffiths and J. Harris. *Principles of Algebraic Geometry*. John Wiley and Sons, Inc., New York, 1994.
- [10] A. Grothendieck and J. Dieudonné. Eléments de géométrie algébrique. *Inst Hautes Études Sci. Publ. Math*, 32, 1967.
- [11] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics, vol.52. Springer Verlag, 1977.
- [12] S. Lang. *Fundamentals of diophantine geometry*. Springer-Verlag, Berlin, 1983.
- [13] S. Lang, editor. *Number Theory III*. Encyclopaedia of Mathematical Sciences, vol.60. Springer-Verlag, Berlin, 1991.
- [14] S. Lang and A. Néron. Rational points of abelian varieties over function fields. *Amer. J. Math.*, 81:95–118, 1959.
- [15] J.S. Milne. *Etale Cohomology*. Princeton University Press, Princeton, NJ, 1980.
- [16] J.S. Milne. *Abelian Varieties*, pages 103–150. Springer-Verlag, Berlin, 1986.
- [17] R. Miranda. Smooth models for elliptic threefolds. In *The Birational Geometry of Degenerations*, Progress in Mathematics, 29. Birkhauser Verlag, 1981.
- [18] R. Miranda. *The Basic Theory of Elliptic Surfaces*. Dottorato di Ricerca in Matematica. ETS Editrice, 1989.
- [19] D. Mumford. *Abelian Varieties*. Tata Institute of Fundamental Research Studies in Mathematics, No. 5. Oxford University Press, 1970.

- [20] D. Mumford. *Geometric Invariant Theory. Second edition.* Ergebnisse der Mathematik und ihrer Grenzgebiete, 34. Springer-Verlag, 1982.
- [21] K. Murty and J. Scherk. Effective versions of the Chebotarev density theorem for function fields. *Comptes Rendus des Seances de l'Academie des Sciences. Serie I. Mathematique*, 319:523–528, 1994.
- [22] K. Nagao. $\mathbb{Q}(t)$ – rank of elliptic curves and certain limits coming from the local points. *Manuscripta Math.*, 92:13–32, 1997.
- [23] J. Neukirch. *Algebraische Zahlentheorie.* Springer Verlag, 1992.
- [24] F. Oort. Algebraic group schemes in characteristic zero are reduced. *Invent. Math.*, 2:79–80, 1966.
- [25] M. Rosen and J. Silverman. On the rank of an elliptic surface. *Invent. Math.*, 133:43–67, 1998.
- [26] J-P. Serre. *Zeta and L functions*, pages 82–92. Harper and Row, New York, 1965.
- [27] T. Shioda. On elliptic modular surfaces. *J. Math. Soc. Japan*, 24:20–59, 1972.
- [28] T. Shioda. Mordell-Weil lattices for higher genus fibration over a curve. In *New trends in algebraic geometry (Warwick, 1996)*, London Math. Soc. Lecture Note Ser., 264, pages 359–373. Cambridge Univ. Press, Cambridge, 1999.
- [29] J. H. Silverman. *Advanced Topics in the Arithmetic of Elliptic Curves.* Graduate Texts in Mathematics, vol.151. Springer Verlag, 1994.
- [30] J. Tate. *Algebraic cycles and the pole of zeta functions*, pages 93–110. Harper and Row, New York, 1965.
- [31] J. Tate. Algorithm for determining the type of a singular fiber in an elliptic pencil. In *Modular Functions of One Variable IV*, Lecture Notes in Mathematics, 476. Springer Verlag, Berlin, 1975.
- [32] R. Wazir. Arithmetic on elliptic threefolds. *Ph.D. thesis*, Brown University, May 2001.